

A/Box Sec

11-29-00

11/28/00  
J0962 U.S. PTO  
09/23/00 4:43:00 PM

# UTILITY PATENT APPLICATION TRANSMITTAL

Submit an original and a duplicate for fee processing  
(Only for new nonprovisional applications under 37 CFR 1.53(b))

## ADDRESS TO:

Assistant Commissioner for Patents  
Box Patent Application  
Washington, D.C. 20231

Attorney Docket No. MBHB00-1214

First Named Inventor Welcher et al

Express Mail No. EL625499212US

Total Pages

US PTO  
09/23/00  
11/28/00

## APPLICATION ELEMENTS

1. ☒ Transmittal Form with Fee
  2. ☒ Specification (including claims and abstract) [Total Pages 118]
  3. ☒ Drawings [Total Sheets 3]
  4. ☐ Oath or Declaration [Total Pages ]
    - a. ☐ Newly executed
    - b. ☐ Copy from prior application

**[Note Boxes 5 and 17 below]**

  - i. ☐ Deletion of Inventor(s) Signed statement attached deleting inventor(s) named in the prior application
5. ☐ Incorporation by Reference: The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
6. ☐ Microfiche Computer Program
7. ☒ Nucleotide and/or Amino Acid Sequence Submission
  - a. ☒ Computer Readable Copy
  - b. ☒ Paper Copy
  - c. ☒ Statement verifying above copies

## ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers
9. ☐ Power of Attorney
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure Statement (IDS)
  - ☐ PTO-1449 Form
  - ☐ Copies of IDS Citations
12. ☐ Preliminary Amendment
13. ☒ Return Receipt Postcard (Should be specifically itemized)
14. ☐ Small Entity Statement(s)
  - ☐ Enclosed
  - ☐ Statement filed in prior application; status still proper and desired
15. ☐ Certified Copy of Priority Document(s)
16. ☒ Other: Title Page

17. ☒ This is a CONTINUING APPLICATION. Please note the following:

- a. ☒ This is a ☒ Continuation ☐ Divisional ☐ Continuation-in-part of prior application Serial No. 60/170,052
- b. ☐ Cancel in this application original claims \_\_\_\_ of the prior application before calculating the filing fee.
- c. ☐ Amend the specification by inserting before the first line the sentence:  
This is a ☐ continuation ☐ divisional ☐ continuation-in-part of application Serial No.
- d. ☒ The prior application is assigned of record to Amgen, Inc.

## UTILITY PATENT APPLICATION TRANSMITTAL

Attorney Docket No. MBHB00-1214

## APPLICATION FEES

BASIC FEE				\$ 710.00
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total Claims	92-20=	72	x \$18.00	\$ 1,296.00
Independent Claims	8-3=	5	x \$80.00	\$ 400.00
<input checked="" type="checkbox"/> Multiple Dependent Claims(s) if applicable			+\$270.00	\$ 270.00
Total of above calculations =				\$ .00
Reduction by 50% for filing by small entity =				\$1/2
<input type="checkbox"/> Assignment fee if applicable			+\$40.00	\$ .00
TOTAL =				\$ 2,676 .00

18. ☐ Please charge my Deposit Account No. 13-2490 in the amount of \$19. ☒ A check in the amount of \$2,676.00 is enclosed.

20. The Commissioner is hereby authorized to credit overpayments or charge any additional fees of the following types to Deposit Account No. 13-2490:

- a. ☒ Fees required under 37 CFR 1.16.
- b. ☒ Fees required under 37 CFR 1.17.
- c. ☒ Fees required under 37 CFR 1.18.

21. ☒ The Commissioner is hereby generally authorized under 37 CFR 1.136(a)(3) to treat any future reply in this or any related application filed pursuant to 37 CFR 1.53 requiring an extension of time as incorporating a request therefor, and the Commissioner is hereby specifically authorized to charge Deposit Account No. 13-2490 for any fee that may be due in connection with such a request for an extension of time.

## 22. CERTIFICATE OF MAILING

I hereby certify that I directed that the correspondence identified above be deposited with the United States Postal Service as "Express Mail Post Office to Addressee" under 37 CFR § 1.10 on the date indicated below and is addressed to the Asst. Commissioner for Patents, Box Patent Application, Washington, DC 20231.

## 23. USPTO CUSTOMER NUMBER

PATENT &amp; TRADEMARK OFFICE



020306

## 24. CORRESPONDENCE ADDRESS

## 25. SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED

Name Reg. No.	Kevin E. Noonan, Reg. No.35,303
Signature	
Date	November 28, 2000

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

Shuqian Jing  
3254 Bordero Lane  
Thousand Oaks, California 91362  
Citizen of the People's Republic of China

Assignee: Amgen, Inc.  
A Corporation of the State of Delaware

Title: Interleukin-1 Receptor Antagonist-Like Molecules and Uses Thereof

# INTERLEUKIN-1 RECEPTOR ANTAGONIST-LIKE MOLECULES AND USES THEREOF

## Field of the Invention

5           The present invention relates to novel Interleukin-1 Receptor Antagonist-Like (IL-1ra-L) polypeptides and nucleic acid molecules encoding the same. The invention also relates to selective binding agents, vectors, host cells, and methods for producing IL-1ra-L polypeptides. The invention further relates to pharmaceutical compositions and methods for the diagnosis, treatment,  
10   amelioration, and/or prevention of diseases, disorders, and conditions associated with IL-1ra-L polypeptides.

## Background of the Invention

15           Technical advances in the identification, cloning, expression, and manipulation of nucleic acid molecules and the deciphering of the human genome have greatly accelerated the discovery of novel therapeutics. Rapid nucleic acid sequencing techniques can now generate sequence information at unprecedented rates and, coupled with computational analyses, allow the assembly of overlapping sequences into partial and entire genomes and the identification of  
20   polypeptide-encoding regions. A comparison of a predicted amino acid sequence against a database compilation of known amino acid sequences allows one to determine the extent of homology to previously identified sequences and/or structural landmarks. The cloning and expression of a polypeptide-encoding region of a nucleic acid molecule provides a polypeptide product for structural  
25   and functional analyses. The manipulation of nucleic acid molecules and encoded polypeptides may confer advantageous properties on a product for use as a therapeutic.

          In spite of the significant technical advances in genome research over the past decade, the potential for the development of novel therapeutics based on the  
30   human genome is still largely unrealized. Many genes encoding potentially beneficial polypeptide therapeutics or those encoding polypeptides, which may



act as “targets” for therapeutic molecules, have still not been identified.

Accordingly, it is an object of the invention to identify novel polypeptides, and nucleic acid molecules encoding the same, which have diagnostic or therapeutic benefit.

5

One of the most potent inflammatory cytokines yet discovered is interleukin-1 (IL-1). IL-1 is thought to be involved in many diseases and medical conditions. It is produced (though not exclusively) by cells of the macrophage/monocyte lineage, and may be produced in two forms: IL-1alpha  
10 (IL-1 $\alpha$ ) and IL-1beta (IL-1 $\beta$ ). Interleukin-1 receptor antagonist (IL-1ra) is a human protein that acts as a natural inhibitor of interleukin-1.

#### Summary of the Invention

The present invention relates to novel IL-1ra-L nucleic acid molecules and  
15 encoded polypeptides.

The invention provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence as set forth in SEQ ID NO: 1;
- (b) the nucleotide sequence of the DNA insert in ATCC Deposit No.  
20 PTA-1215;
- (c) a nucleotide sequence encoding the polypeptide as set forth in SEQ ID NO: 2;
- (d) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (c); and  
25 (e) a nucleotide sequence complementary to any of (a) - (c).

The invention also provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide which is at least  
30 about 70 percent identical to the polypeptide as set forth in SEQ ID NO: 2,

wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(b) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in SEQ ID NO: 1, the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-1215, or (a);

(c) a region of the nucleotide sequence of SEQ ID NO: 1, the DNA insert in ATCC Deposit No. PTA-1215, (a), or (b) encoding a polypeptide fragment of at least about 25 amino acid residues, wherein the polypeptide fragment has an activity of the encoded polypeptide as set forth in SEQ ID NO: 2, or is antigenic;

(d) a region of the nucleotide sequence of SEQ ID NO: 1, the DNA insert in ATCC Deposit No. PTA-1215, or any of (a) - (c) comprising a fragment of at least about 16 nucleotides;

(e) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (d); and

(f) a nucleotide sequence complementary to any of (a) - (d).

The invention further provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one conservative amino acid substitution, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(b) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid insertion, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(c) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid deletion, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(d) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 which has a C- and/or N- terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(e) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(f) a nucleotide sequence of any of (a) - (e) comprising a fragment of at least about 16 nucleotides;

(g) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (f); and

10 (h) a nucleotide sequence complementary to any of (a) - (e).

The present invention provides for an isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence as set forth in SEQ ID NO: 2; and  
15 (b) the amino acid sequence encoded by the DNA insert in ATCC Deposit No. PTA-1215.

The invention also provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

- 20 (a) an amino acid sequence for an ortholog of SEQ ID NO: 2;  
(b) an amino acid sequence which is at least about 70 percent identical to the amino acid sequence of SEQ ID NO: 2, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;  
(c) a fragment of the amino acid sequence set forth in SEQ ID NO: 2  
25 comprising at least about 25 amino acid residues, wherein the fragment has an activity of the polypeptide set forth in SEQ ID NO: 2, or is antigenic; and  
(d) an amino acid sequence for an allelic variant or splice variant of the amino acid sequence as set forth in SEQ ID NO: 2, the amino acid sequence encoded by the DNA insert in ATCC Deposit No. PTA-1215, (a), or (b).

The invention further provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one conservative amino acid substitution, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(b) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one amino acid insertion, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(c) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one amino acid deletion, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(d) the amino acid sequence as set forth in SEQ ID NO: 2 which has a C- and/or N- terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2; and

(e) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2.

Also provided are fusion polypeptides comprising IL-1ra-L amino acid sequences.

The present invention also provides for an expression vector comprising the isolated nucleic acid molecules as set forth herein, recombinant host cells comprising the recombinant nucleic acid molecules as set forth herein, and a method of producing an IL-1ra-L polypeptide comprising culturing the host cells and optionally isolating the polypeptide so produced.

A transgenic non-human animal comprising a nucleic acid molecule encoding an IL-1ra-L polypeptide is also encompassed by the invention. The IL-1ra-L nucleic acid molecules are introduced into the animal in a manner that allows expression and increased levels of an IL-1ra-L polypeptide, which may

include increased circulating levels. Alternatively, the IL-1ra-L nucleic acid molecules are introduced into the animal in a manner that prevents expression of endogenous IL-1ra-L polypeptide (*i.e.*, generates a transgenic animal possessing an IL-1ra-L polypeptide gene knockout). The transgenic non-human animal is  
5 preferably a mammal, and more preferably a rodent, such as a rat or a mouse.

Also provided are derivatives of the IL-1ra-L polypeptides of the present invention.

Additionally provided are selective binding agents such as antibodies and peptides capable of specifically binding the IL-1ra-L polypeptides of the  
10 invention. Such antibodies and peptides may be agonistic or antagonistic.

Pharmaceutical compositions comprising the nucleotides, polypeptides, or selective binding agents of the invention and one or more pharmaceutically acceptable formulation agents are also encompassed by the invention. The pharmaceutical compositions are used to provide therapeutically effective  
15 amounts of the nucleotides or polypeptides of the present invention. The invention is also directed to methods of using the polypeptides, nucleic acid molecules, and selective binding agents.

The IL-1ra-L polypeptides and nucleic acid molecules of the present invention may be used to treat, prevent, ameliorate, and/or detect diseases and  
20 disorders, including those recited herein.

The present invention also provides a method of assaying test molecules to identify a test molecule that binds to an IL-1ra-L polypeptide. The method comprises contacting an IL-1ra-L polypeptide with a test molecule to determine the extent of binding of the test molecule to the polypeptide. The method further  
25 comprises determining whether such test molecules are agonists or antagonists of an IL-1ra-L polypeptide. The present invention further provides a method of testing the impact of molecules on the expression of IL-1ra-L polypeptide or on the activity of IL-1ra-L polypeptide.

Methods of regulating expression and modulating (*i.e.*, increasing or  
30 decreasing) levels of an IL-1ra-L polypeptide are also encompassed by the invention. One method comprises administering to an animal a nucleic acid

molecule encoding an IL-1ra-L polypeptide. In another method, a nucleic acid molecule comprising elements that regulate or modulate the expression of an IL-1ra-L polypeptide may be administered. Examples of these methods include gene therapy, cell therapy, and anti-sense therapy as further described herein.

5 In another aspect of the present invention, the IL-1ra-L polypeptides may be used for identifying receptors thereof ("IL-1ra-L polypeptide receptors"). Various forms of "expression cloning" have been extensively used to clone receptors for protein ligands. *See, e.g.,* Simonsen and Lodish, 1994, *Trends Pharmacol. Sci.* 15:437-41 and Tartaglia *et al.*, 1995, *Cell* 83:1263-71. The  
10 isolation of an IL-1ra-L polypeptide receptor is useful for identifying or developing novel agonists and antagonists of the IL-1ra-L polypeptide signaling pathway. Such agonists and antagonists include soluble IL-1ra-L polypeptide receptors, anti-IL-1ra-L polypeptide receptor-selective binding agents (such as antibodies and derivatives thereof), small molecules, and antisense  
15 oligonucleotides, any of which can be used for treating one or more disease or disorder, including those disclosed herein.

#### Brief Description of the Figures

Figures 1A-1B illustrate the nucleotide sequence of the human IL-1ra-L gene  
20 (SEQ ID NO: 1) and the deduced amino acid sequence of human IL-1ra-L polypeptide (SEQ ID NO: 2);

Figure 2 illustrates the amino acid sequence alignment of human IL-1 $\delta$  (IL-1\_delta; SEQ ID NO: 3), human IL-1ra-L polypeptide (IL-1ra-L; SEQ ID NO: 2),  
25 human IL-1 $\epsilon$  (IL-1\_epsilon; SEQ ID NO: 4), human IL-1 receptor antagonist, secreted polypeptide (IL-1ra\_sec; SEQ ID NO: 9), human IL-1 $\beta$  (IL-1\_beta; SEQ ID NO: 6), and amino acid positions sharing some similarity (consensus).

#### Detailed Description of the Invention

The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All references cited in this application are expressly incorporated by reference herein.

5    Definitions

          The terms “IL-1ra-L gene” or “IL-1ra-L nucleic acid molecule” or “IL-1ra-L polynucleotide” refer to a nucleic acid molecule comprising or consisting of a nucleotide sequence as set forth in SEQ ID NO: 1, a nucleotide sequence encoding the polypeptide as set forth in SEQ ID NO: 2, a nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-1215, and nucleic acid molecules as defined herein.

          The term “IL-1ra-L polypeptide allelic variant” refers to one of several possible naturally occurring alternate forms of a gene occupying a given locus on a chromosome of an organism or a population of organisms.

15       The term “IL-1ra-L polypeptide splice variant” refers to a nucleic acid molecule, usually RNA, which is generated by alternative processing of intron sequences in an RNA transcript of IL-1ra-L polypeptide amino acid sequence as set forth in SEQ ID NO: 2.

          The term “isolated nucleic acid molecule” refers to a nucleic acid molecule of the invention that (1) has been separated from at least about 50 percent of proteins, lipids, carbohydrates, or other materials with which it is naturally found when total nucleic acid is isolated from the source cells, (2) is not linked to all or a portion of a polynucleotide to which the “isolated nucleic acid molecule” is linked in nature, (3) is operably linked to a polynucleotide which it is not linked to in nature, or (4) does not occur in nature as part of a larger polynucleotide sequence. Preferably, the isolated nucleic acid molecule of the present invention is substantially free from any other contaminating nucleic acid molecule(s) or other contaminants that are found in its natural environment that would interfere with its use in polypeptide production or its therapeutic, diagnostic, prophylactic or research use.

The term “nucleic acid sequence” or “nucleic acid molecule” refers to a DNA or RNA sequence. The term encompasses molecules formed from any of the known base analogs of DNA and RNA such as, but not limited to 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinyl-cytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl) uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, N6-iso-pentenyladenine, 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 1-methylinosine, 2,2-dimethyl-guanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyamino-methyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonyl-methyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

The term “vector” is used to refer to any molecule (*e.g.*, nucleic acid, plasmid, or virus) used to transfer coding information to a host cell.

The term “expression vector” refers to a vector that is suitable for transformation of a host cell and contains nucleic acid sequences that direct and/or control the expression of inserted heterologous nucleic acid sequences. Expression includes, but is not limited to, processes such as transcription, translation, and RNA splicing, if introns are present.

The term “operably linked” is used herein to refer to an arrangement of flanking sequences wherein the flanking sequences so described are configured or assembled so as to perform their usual function. Thus, a flanking sequence operably linked to a coding sequence may be capable of effecting the replication, transcription and/or translation of the coding sequence. For example, a coding sequence is operably linked to a promoter when the promoter is capable of directing transcription of that coding sequence. A flanking sequence need not be



contiguous with the coding sequence, so long as it functions correctly. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered “operably linked” to the coding sequence.

5           The term “host cell” is used to refer to a cell which has been transformed, or is capable of being transformed with a nucleic acid sequence and then of expressing a selected gene of interest. The term includes the progeny of the parent cell, whether or not the progeny is identical in morphology or in genetic make-up to the original parent, so long as the selected gene is present.

10           The term “IL-1ra-L polypeptide” refers to a polypeptide comprising the amino acid sequence of SEQ ID NO: 2 and related polypeptides. Related polypeptides include IL-1ra-L polypeptide fragments, IL-1ra-L polypeptide orthologs, IL-1ra-L polypeptide variants, and IL-1ra-L polypeptide derivatives, which possess at least one activity of the polypeptide as set forth in SEQ ID NO:  
15   2. IL-1ra-L polypeptides may be mature polypeptides, as defined herein, and may or may not have an amino-terminal methionine residue, depending on the method by which they are prepared.

          The term “IL-1ra-L polypeptide fragment” refers to a polypeptide that comprises a truncation at the amino-terminus (with or without a leader sequence)  
20   and/or a truncation at the carboxyl-terminus of the polypeptide as set forth in SEQ ID NO: 2. The term “IL-1ra-L polypeptide fragment” also refers to amino-terminal and/or carboxyl-terminal truncations of IL-1ra-L polypeptide orthologs, IL-1ra-L polypeptide derivatives, or IL-1ra-L polypeptide variants, or to amino-terminal and/or carboxyl-terminal truncations of the polypeptides encoded by IL-  
25   1ra-L polypeptide allelic variants or IL-1ra-L polypeptide splice variants. IL-1ra-L polypeptide fragments may result from alternative RNA splicing or from *in vivo* protease activity. Membrane-bound forms of an IL-1ra-L polypeptide are also contemplated by the present invention. In preferred embodiments, truncations and/or deletions comprise about 10 amino acids, or about 20 amino acids, or about  
30   50 amino acids, or about 75 amino acids, or about 100 amino acids, or more than about 100 amino acids. The polypeptide fragments so produced will comprise

about 25 contiguous amino acids, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or about 125 amino acids. Such IL-1ra-L polypeptide fragments may optionally comprise an amino-terminal methionine residue. It will be appreciated that such fragments can be used, for example, to generate antibodies to IL-1ra-L polypeptides.

The term "IL-1ra-L polypeptide ortholog" refers to a polypeptide from another species that corresponds to IL-1ra-L polypeptide amino acid sequence as set forth in SEQ ID NO: 2. For example, mouse and human IL-1ra-L polypeptides are considered orthologs of each other.

The term "IL-1ra-L polypeptide variants" refers to IL-1ra-L polypeptides comprising amino acid sequences having one or more amino acid sequence substitutions, deletions (such as internal deletions and/or IL-1ra-L polypeptide fragments), and/or additions (such as internal additions and/or IL-1ra-L fusion polypeptides) as compared to the IL-1ra-L polypeptide amino acid sequence set forth in SEQ ID NO: 2 (with or without a leader sequence). Variants may be naturally occurring (e.g., IL-1ra-L polypeptide allelic variants, IL-1ra-L polypeptide orthologs, and IL-1ra-L polypeptide splice variants) or artificially constructed. Such IL-1ra-L polypeptide variants may be prepared from the corresponding nucleic acid molecules having a DNA sequence that varies accordingly from the DNA sequence as set forth in SEQ ID NO: 1. In preferred embodiments, the variants have from 1 to 3, or from 1 to 5, or from 1 to 10, or from 1 to 15, or from 1 to 20, or from 1 to 25, or from 1 to 50, or from 1 to 75, or from 1 to 100, or more than 100 amino acid substitutions, insertions, additions and/or deletions, wherein the substitutions may be conservative, or non-conservative, or any combination thereof.

The term "IL-1ra-L polypeptide derivatives" refers to the polypeptide as set forth in SEQ ID NO: 2, IL-1ra-L polypeptide fragments, IL-1ra-L polypeptide orthologs, or IL-1ra-L polypeptide variants, as defined herein, that have been chemically modified. The term "IL-1ra-L polypeptide derivatives" also refers to the polypeptides encoded by IL-1ra-L polypeptide allelic variants or IL-1ra-L

polypeptide splice variants, as defined herein, that have been chemically modified.

The term “mature IL-1ra-L polypeptide” refers to an IL-1ra-L polypeptide lacking a leader sequence. A mature IL-1ra-L polypeptide may also include other  
5 modifications such as proteolytic processing of the amino-terminus (with or without a leader sequence) and/or the carboxyl-terminus, cleavage of a smaller polypeptide from a larger precursor, N-linked and/or O-linked glycosylation, and the like.

The term “IL-1ra-L fusion polypeptide” refers to a fusion of one or more  
10 amino acids (such as a heterologous protein or peptide) at the amino- or carboxyl-terminus of the polypeptide as set forth in SEQ ID NO: 2, IL-1ra-L polypeptide fragments, IL-1ra-L polypeptide orthologs, IL-1ra-L polypeptide variants, or IL-1ra-L derivatives, as defined herein. The term “IL-1ra-L fusion polypeptide” also refers to a fusion of one or more amino acids at the amino- or carboxyl-terminus  
15 of the polypeptide encoded by IL-1ra-L polypeptide allelic variants or IL-1ra-L polypeptide splice variants, as defined herein.

The term “biologically active IL-1ra-L polypeptides” refers to IL-1ra-L polypeptides having at least one activity characteristic of the polypeptide comprising the amino acid sequence of SEQ ID NO: 2. In addition, an IL-1ra-L  
20 polypeptide may be active as an immunogen; that is, the IL-1ra-L polypeptide contains at least one epitope to which antibodies may be raised.

The term “isolated polypeptide” refers to a polypeptide of the present invention that (1) has been separated from at least about 50 percent of polynucleotides, lipids, carbohydrates, or other materials with which it is naturally  
25 found when isolated from the source cell, (2) is not linked (by covalent or noncovalent interaction) to all or a portion of a polypeptide to which the “isolated polypeptide” is linked in nature, (3) is operably linked (by covalent or noncovalent interaction) to a polypeptide with which it is not linked in nature, or (4) does not occur in nature. Preferably, the isolated polypeptide is substantially  
30 free from any other contaminating polypeptides or other contaminants that are found in its natural environment that would interfere with its therapeutic,

diagnostic, prophylactic or research use.

The term "identity," as known in the art, refers to a relationship between the sequences of two or more polypeptide molecules or two or more nucleic acid molecules, as determined by comparing the sequences. In the art, "identity" also  
5 means the degree of sequence relatedness between nucleic acid molecules or polypeptides, as the case may be, as determined by the match between strings of two or more nucleotide or two or more amino acid sequences. "Identity" measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical  
10 model or computer program (*i.e.*, "algorithms").

The term "similarity" is a related concept, but in contrast to "identity," "similarity" refers to a measure of relatedness which includes both identical matches and conservative substitution matches. If two polypeptide sequences have, for example, 10/20 identical amino acids, and the remainder are all non-  
15 conservative substitutions, then the percent identity and similarity would both be 50%. If in the same example, there are five more positions where there are conservative substitutions, then the percent identity remains 50%, but the percent similarity would be 75% (15/20). Therefore, in cases where there are conservative substitutions, the percent similarity between two polypeptides will be  
20 higher than the percent identity between those two polypeptides.

The term "naturally occurring" or "native" when used in connection with biological materials such as nucleic acid molecules, polypeptides, host cells, and the like, refers to materials which are found in nature and are not manipulated by man. Similarly, "non-naturally occurring" or "non-native" as used herein refers to  
25 a material that is not found in nature or that has been structurally modified or synthesized by man.

The terms "effective amount" and "therapeutically effective amount" each refer to the amount of an IL-1ra-L polypeptide or IL-1ra-L nucleic acid molecule used to support an observable level of one or more biological activities of the IL-  
30 1ra-L polypeptides as set forth herein.

The term “pharmaceutically acceptable carrier” or “physiologically acceptable carrier” as used herein refers to one or more formulation materials suitable for accomplishing or enhancing the delivery of the IL-1ra-L polypeptide, IL-1ra-L nucleic acid molecule, or IL-1ra-L selective binding agent as a pharmaceutical composition.

The term “antigen” refers to a molecule or a portion of a molecule capable of being bound by a selective binding agent, such as an antibody, and additionally capable of being used in an animal to produce antibodies capable of binding to an epitope of that antigen. An antigen may have one or more epitopes.

The term “selective binding agent” refers to a molecule or molecules having specificity for an IL-1ra-L polypeptide. As used herein, the terms, “specific” and “specificity” refer to the ability of the selective binding agents to bind to human IL-1ra-L polypeptides and not to bind to human non-IL-1ra-L polypeptides. It will be appreciated, however, that the selective binding agents may also bind orthologs of the polypeptide as set forth in SEQ ID NO: 2, that is, interspecies versions thereof, such as mouse and rat IL-1ra-L polypeptides.

The term “transduction” is used to refer to the transfer of genes from one bacterium to another, usually by a phage. “Transduction” also refers to the acquisition and transfer of eukaryotic cellular sequences by retroviruses.

The term “transfection” is used to refer to the uptake of foreign or exogenous DNA by a cell, and a cell has been “transfected” when the exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are well known in the art and are disclosed herein. *See, e.g.,* Graham *et al.*, 1973, *Virology* 52:456; Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratories, 1989); Davis *et al.*, *Basic Methods in Molecular Biology* (Elsevier, 1986); and Chu *et al.*, 1981, *Gene* 13:197. Such techniques can be used to introduce one or more exogenous DNA moieties into suitable host cells.

The term “transformation” as used herein refers to a change in a cell’s genetic characteristics, and a cell has been transformed when it has been modified to contain a new DNA. For example, a cell is transformed where it is genetically

modified from its native state. Following transfection or transduction, the transforming DNA may recombine with that of the cell by physically integrating into a chromosome of the cell, may be maintained transiently as an episomal element without being replicated, or may replicate independently as a plasmid. A  
5 cell is considered to have been stably transformed when the DNA is replicated with the division of the cell.

#### Relatedness of Nucleic Acid Molecules and/or Polypeptides

It is understood that related nucleic acid molecules include allelic or splice  
10 variants of the nucleic acid molecule of SEQ ID NO: 1, and include sequences which are complementary to any of the above nucleotide sequences. Related nucleic acid molecules also include a nucleotide sequence encoding a polypeptide comprising or consisting essentially of a substitution, modification, addition and/or deletion of one or more amino acid residues compared to the polypeptide  
15 in SEQ ID NO: 2. Such related IL-1ra-L polypeptides may comprise, for example, an addition and/or a deletion of one or more N-linked or O-linked glycosylation sites or an addition and/or a deletion of one or more cysteine residues.

Related nucleic acid molecules also include fragments of IL-1ra-L nucleic  
20 acid molecules which encode a polypeptide of at least about 25 contiguous amino acids, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or about 125 amino acids, or more than 125 amino acid residues of the IL-1ra-L polypeptide of SEQ ID NO: 2.

In addition, related IL-1ra-L nucleic acid molecules also include those  
25 molecules which comprise nucleotide sequences which hybridize under moderately or highly stringent conditions as defined herein with the fully complementary sequence of the IL-1ra-L nucleic acid molecule of SEQ ID NO: 1, or of a molecule encoding a polypeptide, which polypeptide comprises the amino acid sequence as shown in SEQ ID NO: 2, or of a nucleic acid fragment as  
30 defined herein, or of a nucleic acid fragment encoding a polypeptide as defined herein. Hybridization probes may be prepared using the IL-1ra-L sequences

provided herein to screen cDNA, genomic or synthetic DNA libraries for related sequences. Regions of the DNA and/or amino acid sequence of IL-1ra-L polypeptide that exhibit significant identity to known sequences are readily determined using sequence alignment algorithms as described herein and those  
5 regions may be used to design probes for screening.

The term "highly stringent conditions" refers to those conditions that are designed to permit hybridization of DNA strands whose sequences are highly complementary, and to exclude hybridization of significantly mismatched DNAs. Hybridization stringency is principally determined by temperature, ionic strength,  
10 and the concentration of denaturing agents such as formamide. Examples of "highly stringent conditions" for hybridization and washing are 0.015 M sodium chloride, 0.0015 M sodium citrate at 65-68°C or 0.015 M sodium chloride, 0.0015 M sodium citrate, and 50% formamide at 42°C. See Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual* (2nd ed., Cold Spring Harbor  
15 Laboratory, 1989); Anderson *et al.*, *Nucleic Acid Hybridisation: A Practical Approach* Ch. 4 (IRL Press Limited).

More stringent conditions (such as higher temperature, lower ionic strength, higher formamide, or other denaturing agent) may also be used – however, the rate of hybridization will be affected. Other agents may be included  
20 in the hybridization and washing buffers for the purpose of reducing non-specific and/or background hybridization. Examples are 0.1% bovine serum albumin, 0.1% polyvinyl-pyrrolidone, 0.1% sodium pyrophosphate, 0.1% sodium dodecylsulfate, NaDodSO<sub>4</sub>, (SDS), ficoll, Denhardt's solution, sonicated salmon sperm DNA (or another non-complementary DNA), and dextran sulfate, although  
25 other suitable agents can also be used. The concentration and types of these additives can be changed without substantially affecting the stringency of the hybridization conditions. Hybridization experiments are usually carried out at pH 6.8-7.4; however, at typical ionic strength conditions, the rate of hybridization is nearly independent of pH. See Anderson *et al.*, *Nucleic Acid Hybridisation: A  
30 Practical Approach* Ch. 4 (IRL Press Limited).

Factors affecting the stability of DNA duplex include base composition, length, and degree of base pair mismatch. Hybridization conditions can be adjusted by one skilled in the art in order to accommodate these variables and allow DNAs of different sequence relatedness to form hybrids. The melting  
5 temperature of a perfectly matched DNA duplex can be estimated by the following equation:

$$T_m(^{\circ}\text{C}) = 81.5 + 16.6(\log[\text{Na}^+]) + 0.41(\% \text{G+C}) - 600/N - 0.72(\% \text{formamide})$$

where N is the length of the duplex formed, [Na<sup>+</sup>] is the molar concentration of the sodium ion in the hybridization or washing solution, %G+C is the percentage  
10 of (guanine+cytosine) bases in the hybrid. For imperfectly matched hybrids, the melting temperature is reduced by approximately 1°C for each 1% mismatch.

The term “moderately stringent conditions” refers to conditions under which a DNA duplex with a greater degree of base pair mismatching than could occur under “highly stringent conditions” is able to form. Examples of typical  
15 “moderately stringent conditions” are 0.015 M sodium chloride, 0.0015 M sodium citrate at 50-65°C or 0.015 M sodium chloride, 0.0015 M sodium citrate, and 20% formamide at 37-50°C. By way of example, “moderately stringent conditions” of 50°C in 0.015 M sodium ion will allow about a 21% mismatch.

It will be appreciated by those skilled in the art that there is no absolute  
20 distinction between “highly stringent conditions” and “moderately stringent conditions.” For example, at 0.015 M sodium ion (no formamide), the melting temperature of perfectly matched long DNA is about 71°C. With a wash at 65°C (at the same ionic strength), this would allow for approximately a 6% mismatch. To capture more distantly related sequences, one skilled in the art can simply  
25 lower the temperature or raise the ionic strength.

A good estimate of the melting temperature in 1M NaCl\* for oligonucleotide probes up to about 20nt is given by:

$$T_m = 2^{\circ}\text{C per A-T base pair} + 4^{\circ}\text{C per G-C base pair}$$

\*The sodium ion concentration in 6X salt sodium citrate (SSC) is 1M. *See Suggs et al., Developmental Biology Using Purified Genes* 683 (Brown and Fox, eds.,  
30 1981).



High stringency washing conditions for oligonucleotides are usually at a temperature of 0-5°C below the T<sub>m</sub> of the oligonucleotide in 6X SSC, 0.1% SDS.

In another embodiment, related nucleic acid molecules comprise or consist of a nucleotide sequence that is at least about 70 percent identical to the nucleotide sequence as shown in SEQ ID NO: 1, or comprise or consist essentially of a nucleotide sequence encoding a polypeptide that is at least about 70 percent identical to the polypeptide as set forth in SEQ ID NO: 2. In preferred embodiments, the nucleotide sequences are about 75 percent, or about 80 percent, or about 85 percent, or about 90 percent, or about 95, 96, 97, 98, or 99 percent identical to the nucleotide sequence as shown in SEQ ID NO: 1, or the nucleotide sequences encode a polypeptide that is about 75 percent, or about 80 percent, or about 85 percent, or about 90 percent, or about 95, 96, 97, 98, or 99 percent identical to the polypeptide sequence as set forth in SEQ ID NO: 2. Related nucleic acid molecules encode polypeptides possessing at least one activity of the polypeptide set forth in SEQ ID NO: 2.

Differences in the nucleic acid sequence may result in conservative and/or non-conservative modifications of the amino acid sequence relative to the amino acid sequence of in SEQ ID NO: 2.

Conservative modifications to the amino acid sequence of SEQ ID NO: 2 (and the corresponding modifications to the encoding nucleotides) will produce a polypeptide having functional and chemical characteristics similar to those of IL-1ra-L polypeptides. In contrast, substantial modifications in the functional and/or chemical characteristics of IL-1ra-L polypeptides may be accomplished by selecting substitutions in the amino acid sequence of SEQ ID NO: 2 that differ significantly in their effect on maintaining (a) the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

For example, a “conservative amino acid substitution” may involve a substitution of a native amino acid residue with a nonnative residue such that there is little or no effect on the polarity or charge of the amino acid residue at

that position. Furthermore, any native residue in the polypeptide may also be substituted with alanine, as has been previously described for “alanine scanning mutagenesis.”

Conservative amino acid substitutions also encompass non-naturally occurring amino acid residues that are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics, and other reversed or inverted forms of amino acid moieties.

Naturally occurring residues may be divided into classes based on common side chain properties:

- 1) hydrophobic: norleucine, Met, Ala, Val, Leu, Ile;
- 2) neutral hydrophilic: Cys, Ser, Thr;
- 3) acidic: Asp, Glu;
- 4) basic: Asn, Gln, His, Lys, Arg;
- 5) residues that influence chain orientation: Gly, Pro; and
- 6) aromatic: Trp, Tyr, Phe.

For example, non-conservative substitutions may involve the exchange of a member of one of these classes for a member from another class. Such substituted residues may be introduced into regions of the human IL-1ra-L polypeptide that are homologous with non-human IL-1ra-L polypeptides, or into the non-homologous regions of the molecule.

In making such changes, the hydropathic index of amino acids may be considered. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics. The hydropathic indices are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

The importance of the hydropathic amino acid index in conferring interactive biological function on a protein is generally understood in the art (Kyte *et al.*, 1982, *J. Mol. Biol.* 157:105-31). It is known that certain amino acids may

be substituted for other amino acids having a similar hydropathic index or score and still retain a similar biological activity. In making changes based upon the hydropathic index, the substitution of amino acids whose hydropathic indices are within  $\pm 2$  is preferred, those which are within  $\pm 1$  are particularly preferred, and  
5 those within  $\pm 0.5$  are even more particularly preferred.

It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity, particularly where the biologically functionally equivalent protein or peptide thereby created is intended for use in immunological embodiments, as in the present case. The greatest local  
10 average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigenicity, *i.e.*, with a biological property of the protein.

The following hydrophilicity values have been assigned to these amino acid residues: arginine (+3.0); lysine (+3.0); aspartate ( $+3.0 \pm 1$ ); glutamate ( $+3.0 \pm 1$ ); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline ( $-0.5 \pm 1$ ); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); and tryptophan (-3.4). In making changes based upon  
15 similar hydrophilicity values, the substitution of amino acids whose hydrophilicity values are within  $\pm 2$  is preferred, those which are within  $\pm 1$  are particularly preferred, and those within  $\pm 0.5$  are even more particularly preferred. One may also identify epitopes from primary amino acid sequences on the basis of hydrophilicity. These regions are also referred to as "epitopic core regions."

Desired amino acid substitutions (whether conservative or non-conservative) can be determined by those skilled in the art at the time such  
25 substitutions are desired. For example, amino acid substitutions can be used to identify important residues of the IL-1ra-L polypeptide, or to increase or decrease the affinity of the IL-1ra-L polypeptides described herein. Exemplary amino acid substitutions are set forth in Table I.

Amino Acid Substitutions

Original Residues	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln	Gln
Asp	Glu	Glu
Cys	Ser, Ala	Ser
Gln	Asn	Asn
Glu	Asp	Asp
Gly	Pro, Ala	Ala
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe, Norleucine	Leu
Leu	Norleucine, Ile, Val, Met, Ala, Phe	Ile
Lys	Arg, 1,4 Diamino-butyric Acid, Gln, Asn	Arg
Met	Leu, Phe, Ile	Leu
Phe	Leu, Val, Ile, Ala, Tyr	Leu
Pro	Ala	Gly
Ser	Thr, Ala, Cys	Thr
Thr	Ser	Ser
Trp	Tyr, Phe	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Met, Leu, Phe, Ala, Norleucine	Leu

A skilled artisan will be able to determine suitable variants of the polypeptide as set forth in SEQ ID NO: 2 using well-known techniques. For

identifying suitable areas of the molecule that may be changed without destroying biological activity, one skilled in the art may target areas not believed to be important for activity. For example, when similar polypeptides with similar activities from the same species or from other species are known, one skilled in the art may compare the amino acid sequence of an IL-1ra-L polypeptide to such similar polypeptides. With such a comparison, one can identify residues and portions of the molecules that are conserved among similar polypeptides. It will be appreciated that changes in areas of the IL-1ra-L molecule that are not conserved relative to such similar polypeptides would be less likely to adversely affect the biological activity and/or structure of an IL-1ra-L polypeptide. One skilled in the art would also know that, even in relatively conserved regions, one may substitute chemically similar amino acids for the naturally occurring residues while retaining activity (conservative amino acid residue substitutions). Therefore, even areas that may be important for biological activity or for structure may be subject to conservative amino acid substitutions without destroying the biological activity or without adversely affecting the polypeptide structure.

Additionally, one skilled in the art can review structure-function studies identifying residues in similar polypeptides that are important for activity or structure. In view of such a comparison, one can predict the importance of amino acid residues in an IL-1ra-L polypeptide that correspond to amino acid residues that are important for activity or structure in similar polypeptides. One skilled in the art may opt for chemically similar amino acid substitutions for such predicted important amino acid residues of IL-1ra-L polypeptides.

One skilled in the art can also analyze the three-dimensional structure and amino acid sequence in relation to that structure in similar polypeptides. In view of such information, one skilled in the art may predict the alignment of amino acid residues of IL-1ra-L polypeptide with respect to its three dimensional structure. One skilled in the art may choose not to make radical changes to amino acid residues predicted to be on the surface of the protein, since such residues may be involved in important interactions with other molecules. Moreover, one skilled in the art may generate test variants containing a single amino acid substitution at

each amino acid residue. The variants could be screened using activity assays known to those with skill in the art. Such variants could be used to gather information about suitable variants. For example, if one discovered that a change to a particular amino acid residue resulted in destroyed, undesirably reduced, or unsuitable activity, variants with such a change would be avoided. In other words, based on information gathered from such routine experiments, one skilled in the art can readily determine the amino acids where further substitutions should be avoided either alone or in combination with other mutations.

A number of scientific publications have been devoted to the prediction of secondary structure. See Moulton, 1996, *Curr. Opin. Biotechnol.* 7:422-27; Chou *et al.*, 1974, *Biochemistry* 13:222-45; Chou *et al.*, 1974, *Biochemistry* 113:211-22; Chou *et al.*, 1978, *Adv. Enzymol. Relat. Areas Mol. Biol.* 47:45-48; Chou *et al.*, 1978, *Ann. Rev. Biochem.* 47:251-276; and Chou *et al.*, 1979, *Biophys. J.* 26:367-84. Moreover, computer programs are currently available to assist with predicting secondary structure. One method of predicting secondary structure is based upon homology modeling. For example, two polypeptides or proteins which have a sequence identity of greater than 30%, or similarity greater than 40%, often have similar structural topologies. The recent growth of the protein structural database (PDB) has provided enhanced predictability of secondary structure, including the potential number of folds within the structure of a polypeptide or protein. See Holm *et al.*, 1999, *Nucleic Acids Res.* 27:244-47. It has been suggested that there are a limited number of folds in a given polypeptide or protein and that once a critical number of structures have been resolved, structural prediction will become dramatically more accurate (Brenner *et al.*, 1997, *Curr. Opin. Struct. Biol.* 7:369-76).

Additional methods of predicting secondary structure include "threading" (Jones, 1997, *Curr. Opin. Struct. Biol.* 7:377-87; Sippl *et al.*, 1996, *Structure* 4:15-19), "profile analysis" (Bowie *et al.*, 1991, *Science*, 253:164-70; Gribskov *et al.*, 1990, *Methods Enzymol.* 183:146-59; Gribskov *et al.*, 1987, *Proc. Nat. Acad. Sci. U.S.A.* 84:4355-58), and "evolutionary linkage" (See Holm *et al.*, *supra*, and Brenner *et al.*, *supra*).

Preferred IL-1ra-L polypeptide variants include glycosylation variants wherein the number and/or type of glycosylation sites have been altered compared to the amino acid sequence set forth in SEQ ID NO: 2. In one embodiment, IL-1ra-L polypeptide variants comprise a greater or a lesser number of N-linked glycosylation sites than the amino acid sequence set forth in SEQ ID NO: 2. An N-linked glycosylation site is characterized by the sequence: Asn-X-Ser or Asn-X-Thr, wherein the amino acid residue designated as X may be any amino acid residue except proline. The substitution of amino acid residues to create this sequence provides a potential new site for the addition of an N-linked carbohydrate chain. Alternatively, substitutions that eliminate this sequence will remove an existing N-linked carbohydrate chain. Also provided is a rearrangement of N-linked carbohydrate chains wherein one or more N-linked glycosylation sites (typically those that are naturally occurring) are eliminated and one or more new N-linked sites are created. Additional preferred IL-1ra-L variants include cysteine variants, wherein one or more cysteine residues are deleted or substituted with another amino acid (*e.g.*, serine) as compared to the amino acid sequence set forth in SEQ ID NO: 2. Cysteine variants are useful when IL-1ra-L polypeptides must be refolded into a biologically active conformation such as after the isolation of insoluble inclusion bodies. Cysteine variants generally have fewer cysteine residues than the native protein, and typically have an even number to minimize interactions resulting from unpaired cysteines.

In other embodiments, related nucleic acid molecules comprise or consist of a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid insertion and wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2, or a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid deletion and wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2. Related nucleic acid molecules also comprise or consist of a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 wherein the polypeptide has a carboxyl- and/or amino-terminal truncation and further wherein

the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2. Related nucleic acid molecules also comprise or consist of a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, carboxyl-terminal truncations, and amino-terminal truncations and wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2.

In addition, the polypeptide comprising the amino acid sequence of SEQ ID NO: 2, or other IL-1ra-L polypeptide, may be fused to a homologous polypeptide to form a homodimer or to a heterologous polypeptide to form a heterodimer. Heterologous peptides and polypeptides include, but are not limited to: an epitope to allow for the detection and/or isolation of an IL-1ra-L fusion polypeptide; a transmembrane receptor protein or a portion thereof, such as an extracellular domain or a transmembrane and intracellular domain; a ligand or a portion thereof which binds to a transmembrane receptor protein; an enzyme or portion thereof which is catalytically active; a polypeptide or peptide which promotes oligomerization, such as a leucine zipper domain; a polypeptide or peptide which increases stability, such as an immunoglobulin constant region; and a polypeptide which has a therapeutic activity different from the polypeptide comprising the amino acid sequence as set forth in SEQ ID NO: 2, or other IL-1ra-L polypeptide.

Fusions can be made either at the amino-terminus or at the carboxyl-terminus of the polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 2, or other IL-1ra-L polypeptide. Fusions may be direct with no linker or adapter molecule or may be through a linker or adapter molecule. A linker or adapter molecule may be one or more amino acid residues, typically from about 20 to about 50 amino acid residues. A linker or adapter molecule may also be designed with a cleavage site for a DNA restriction endonuclease or for a protease to allow for the separation of the fused moieties. It will be appreciated that once constructed, the fusion polypeptides can be derivatized according to the methods described herein.



In a further embodiment of the invention, the polypeptide comprising the amino acid sequence of SEQ ID NO: 2, or other IL-1ra-L polypeptide, is fused to one or more domains of an Fc region of human IgG. Antibodies comprise two functionally independent parts, a variable domain known as "Fab," that binds an antigen, and a constant domain known as "Fc," that is involved in effector functions such as complement activation and attack by phagocytic cells. An Fc has a long serum half-life, whereas an Fab is short-lived. Capon *et al.*, 1989, *Nature* 337:525-31. When constructed together with a therapeutic protein, an Fc domain can provide longer half-life or incorporate such functions as Fc receptor binding, protein A binding, complement fixation, and perhaps even placental transfer. *Id.* Table II summarizes the use of certain Fc fusions known in the art.

Table II  
Fc Fusion with Therapeutic Proteins

Form of Fc	Fusion partner	Therapeutic implications	Reference
IgG1	N-terminus of CD30-L	Hodgkin's disease; anaplastic lymphoma; T-cell leukemia	U.S. Patent No. 5,480,981
Murine Fcγ2a	IL-10	anti-inflammatory; transplant rejection	Zheng <i>et al.</i> , 1995, <i>J. Immunol.</i> 154:5590-600
IgG1	TNF receptor	septic shock	Fisher <i>et al.</i> , 1996, <i>N. Engl. J. Med.</i> 334:1697-1702; Van Zee <i>et al.</i> , 1996, <i>J. Immunol.</i> 156:2221-30
IgG, IgA, IgM, or IgE (excluding the first domain)	TNF receptor	inflammation, autoimmune disorders	U.S. Patent No. 5,808,029
IgG1	CD4 receptor	AIDS	Capon <i>et al.</i> , 1989, <i>Nature</i> 337: 525-31
IgG1, IgG3	N-terminus of IL-2	anti-cancer, antiviral	Harvill <i>et al.</i> , 1995, <i>Immunotech.</i> 1:95-105
IgG1	C-terminus of OPG	osteoarthritis; bone density	WO 97/23614
IgG1	N-terminus of leptin	anti-obesity	PCT/US 97/23183, filed December 11, 1997
Human Ig Cγ1	CTLA-4	autoimmune disorders	Linsley, 1991, <i>J. Exp.</i>

In one example, a human IgG hinge, CH2, and CH3 region may be fused at either the amino-terminus or carboxyl-terminus of the IL-1ra-L polypeptides using methods known to the skilled artisan. In another example, a human IgG hinge, CH2, and CH3 region may be fused at either the amino-terminus or carboxyl-terminus of an IL-1ra-L polypeptide fragment (e.g., the predicted extracellular portion of IL-1ra-L polypeptide).

The resulting IL-1ra-L fusion polypeptide may be purified by use of a Protein A affinity column. Peptides and proteins fused to an Fc region have been found to exhibit a substantially greater half-life *in vivo* than the unfused counterpart. Also, a fusion to an Fc region allows for dimerization/multimerization of the fusion polypeptide. The Fc region may be a naturally occurring Fc region, or may be altered to improve certain qualities, such as therapeutic qualities, circulation time, or reduced aggregation.

Identity and similarity of related nucleic acid molecules and polypeptides are readily calculated by known methods. Such methods include, but are not limited to those described in *Computational Molecular Biology* (A.M. Lesk, ed., Oxford University Press 1988); *Biocomputing: Informatics and Genome Projects* (D.W. Smith, ed., Academic Press 1993); *Computer Analysis of Sequence Data* (Part 1, A.M. Griffin and H.G. Griffin, eds., Humana Press 1994); G. von Heinle, *Sequence Analysis in Molecular Biology* (Academic Press 1987); *Sequence Analysis Primer* (M. Gribskov and J. Devereux, eds., M. Stockton Press 1991); and Carillo *et al.*, 1988, *SIAM J. Applied Math.*, 48:1073.

Preferred methods to determine identity and/or similarity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are described in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package, including GAP (Devereux *et al.*, 1984, *Nucleic Acids Res.* 12:387; Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN,

and FASTA (Altschul *et al.*, 1990, *J. Mol. Biol.* 215:403-10). The BLASTX program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (Altschul *et al.*, *BLAST Manual* (NCB NLM NIH, Bethesda, MD); Altschul *et al.*, 1990, *supra*). The well-known Smith  
5 Waterman algorithm may also be used to determine identity.

Certain alignment schemes for aligning two amino acid sequences may result in the matching of only a short region of the two sequences, and this small aligned region may have very high sequence identity even though there is no significant relationship between the two full-length sequences. Accordingly, in a  
10 preferred embodiment, the selected alignment method (GAP program) will result in an alignment that spans at least 50 contiguous amino acids of the claimed polypeptide.

For example, using the computer algorithm GAP (Genetics Computer Group, University of Wisconsin, Madison, WI), two polypeptides for which the  
15 percent sequence identity is to be determined are aligned for optimal matching of their respective amino acids (the "matched span," as determined by the algorithm). A gap opening penalty (which is calculated as 3X the average diagonal; the "average diagonal" is the average of the diagonal of the comparison matrix being used; the "diagonal" is the score or number assigned to each perfect  
20 amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually 0.1X the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. A standard comparison matrix is also used by the algorithm (*see* Dayhoff *et al.*, 5 *Atlas of Protein Sequence and Structure* (Supp. 3  
25 1978)(PAM250 comparison matrix); Henikoff *et al.*, 1992, *Proc. Natl. Acad. Sci USA* 89:10915-19 (BLOSUM 62 comparison matrix)).

Preferred parameters for polypeptide sequence comparison include the following:

30       Algorithm: Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443-53;  
      Comparison matrix: BLOSUM 62 (Henikoff *et al.*, *supra*);

Gap Penalty: 12

Gap Length Penalty: 4

Threshold of Similarity: 0

- 5 The GAP program is useful with the above parameters. The aforementioned parameters are the default parameters for polypeptide comparisons (along with no penalty for end gaps) using the GAP algorithm.

Preferred parameters for nucleic acid molecule sequence comparison include the following:

10

Algorithm: Needleman and Wunsch, *supra*;

Comparison matrix: matches = +10, mismatch = 0

Gap Penalty: 50

Gap Length Penalty: 3

15

The GAP program is also useful with the above parameters. The aforementioned parameters are the default parameters for nucleic acid molecule comparisons.

- Other exemplary algorithms, gap opening penalties, gap extension penalties, comparison matrices, and thresholds of similarity may be used, including those set forth in the Program Manual, Wisconsin Package, Version 9, September, 1997. The particular choices to be made will be apparent to those of skill in the art and will depend on the specific comparison to be made, such as DNA-to-DNA, protein-to-protein, protein-to-DNA; and additionally, whether the comparison is between given pairs of sequences (in which case GAP or BestFit are generally preferred) or between one sequence and a large database of sequences (in which case FASTA or BLASTA are preferred).
- 20
- 25

#### Nucleic Acid Molecules

- The nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of an IL-1ra-L polypeptide can readily be obtained in a variety of
- 30

ways including, without limitation, chemical synthesis, cDNA or genomic library screening, expression library screening, and/or PCR amplification of cDNA.

Recombinant DNA methods used herein are generally those set forth in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) and/or *Current Protocols in Molecular Biology* (Ausubel *et al.*, eds., Green Publishers Inc. and Wiley and Sons 1994). The invention provides for nucleic acid molecules as described herein and methods for obtaining such molecules.

Where a gene encoding the amino acid sequence of an IL-1ra-L polypeptide has been identified from one species, all or a portion of that gene may be used as a probe to identify orthologs or related genes from the same species. The probes or primers may be used to screen cDNA libraries from various tissue sources believed to express the IL-1ra-L polypeptide. In addition, part or all of a nucleic acid molecule having the sequence as set forth in SEQ ID NO: 1 may be used to screen a genomic library to identify and isolate a gene encoding the amino acid sequence of an IL-1ra-L polypeptide. Typically, conditions of moderate or high stringency will be employed for screening to minimize the number of false positives obtained from the screening.

Nucleic acid molecules encoding the amino acid sequence of IL-1ra-L polypeptides may also be identified by expression cloning which employs the detection of positive clones based upon a property of the expressed protein. Typically, nucleic acid libraries are screened by the binding an antibody or other binding partner (*e.g.*, receptor or ligand) to cloned proteins that are expressed and displayed on a host cell surface. The antibody or binding partner is modified with a detectable label to identify those cells expressing the desired clone.

Recombinant expression techniques conducted in accordance with the descriptions set forth below may be followed to produce these polynucleotides and to express the encoded polypeptides. For example, by inserting a nucleic acid sequence that encodes the amino acid sequence of an IL-1ra-L polypeptide into an appropriate vector, one skilled in the art can readily produce large quantities of the desired nucleotide sequence. The sequences can then be used to generate

detection probes or amplification primers. Alternatively, a polynucleotide encoding the amino acid sequence of an IL-1ra-L polypeptide can be inserted into an expression vector. By introducing the expression vector into an appropriate host, the encoded IL-1ra-L polypeptide may be produced in large amounts.

5 Another method for obtaining a suitable nucleic acid sequence is the polymerase chain reaction (PCR). In this method, cDNA is prepared from poly(A)+RNA or total RNA using the enzyme reverse transcriptase. Two primers, typically complementary to two separate regions of cDNA encoding the amino acid sequence of an IL-1ra-L polypeptide, are then added to the cDNA  
10 along with a polymerase such as *Taq* polymerase, and the polymerase amplifies the cDNA region between the two primers.

Another means of preparing a nucleic acid molecule encoding the amino acid sequence of an IL-1ra-L polypeptide is chemical synthesis using methods well known to the skilled artisan such as those described by Engels *et al.*, 1989,  
15 *Angew. Chem. Intl. Ed.* 28:716-34. These methods include, *inter alia*, the phosphotriester, phosphoramidite, and H-phosphonate methods for nucleic acid synthesis. A preferred method for such chemical synthesis is polymer-supported synthesis using standard phosphoramidite chemistry. Typically, the DNA encoding the amino acid sequence of an IL-1ra-L polypeptide will be several  
20 hundred nucleotides in length. Nucleic acids larger than about 100 nucleotides can be synthesized as several fragments using these methods. The fragments can then be ligated together to form the full-length nucleotide sequence of an IL-1ra-L gene. Usually, the DNA fragment encoding the amino-terminus of the polypeptide will have an ATG, which encodes a methionine residue. This  
25 methionine may or may not be present on the mature form of the IL-1ra-L polypeptide, depending on whether the polypeptide produced in the host cell is designed to be secreted from that cell. Other methods known to the skilled artisan may be used as well.

In certain embodiments, nucleic acid variants contain codons which have  
30 been altered for optimal expression of an IL-1ra-L polypeptide in a given host cell. Particular codon alterations will depend upon the IL-1ra-L polypeptide and

host cell selected for expression. Such "codon optimization" can be carried out by a variety of methods, for example, by selecting codons which are preferred for use in highly expressed genes in a given host cell. Computer algorithms which incorporate codon frequency tables such as "Eco\_high.Cod" for codon preference of highly expressed bacterial genes may be used and are provided by the University of Wisconsin Package Version 9.0 (Genetics Computer Group, Madison, WI). Other useful codon frequency tables include "Celegans\_high.cod," "Celegans\_low.cod," "Drosophila\_high.cod," "Human\_high.cod," "Maize\_high.cod," and "Yeast\_high.cod."

In some cases, it may be desirable to prepare nucleic acid molecules encoding IL-1ra-L polypeptide variants. Nucleic acid molecules encoding variants may be produced using site directed mutagenesis, PCR amplification, or other appropriate methods, where the primer(s) have the desired point mutations (*see* Sambrook *et al.*, *supra*, and Ausubel *et al.*, *supra*, for descriptions of mutagenesis techniques). Chemical synthesis using methods described by Engels *et al.*, *supra*, may also be used to prepare such variants. Other methods known to the skilled artisan may be used as well.

#### Vectors and Host Cells

A nucleic acid molecule encoding the amino acid sequence of an IL-1ra-L polypeptide is inserted into an appropriate expression vector using standard ligation techniques. The vector is typically selected to be functional in the particular host cell employed (*i.e.*, the vector is compatible with the host cell machinery such that amplification of the gene and/or expression of the gene can occur). A nucleic acid molecule encoding the amino acid sequence of an IL-1ra-L polypeptide may be amplified/expressed in prokaryotic, yeast, insect (baculovirus systems) and/or eukaryotic host cells. Selection of the host cell will depend in part on whether an IL-1ra-L polypeptide is to be post-translationally modified (*e.g.*, glycosylated and/or phosphorylated). If so, yeast, insect, or mammalian host cells are preferable. For a review of expression vectors, *see Meth. Enz.*, vol. 185 (D.V. Goeddel, ed., Academic Press 1990).

Typically, expression vectors used in any of the host cells will contain sequences for plasmid maintenance and for cloning and expression of exogenous nucleotide sequences. Such sequences, collectively referred to as “flanking sequences” in certain embodiments will typically include one or more of the following nucleotide sequences: a promoter, one or more enhancer sequences, an origin of replication, a transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a sequence encoding a leader sequence for polypeptide secretion, a ribosome binding site, a polyadenylation sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide to be expressed, and a selectable marker element. Each of these sequences is discussed below.

Optionally, the vector may contain a “tag”-encoding sequence, *i.e.*, an oligonucleotide molecule located at the 5’ or 3’ end of the IL-1ra-L polypeptide coding sequence; the oligonucleotide sequence encodes polyHis (such as hexaHis), or another “tag” such as FLAG, HA (hemagglutinin influenza virus), or *myc* for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for affinity purification of the IL-1ra-L polypeptide from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can subsequently be removed from the purified IL-1ra-L polypeptide by various means such as using certain peptidases for cleavage.

Flanking sequences may be homologous (*i.e.*, from the same species and/or strain as the host cell), heterologous (*i.e.*, from a species other than the host cell species or strain), hybrid (*i.e.*, a combination of flanking sequences from more than one source), or synthetic, or the flanking sequences may be native sequences which normally function to regulate IL-1ra-L polypeptide expression. As such, the source of a flanking sequence may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence is functional in, and can be activated by, the host cell machinery.



Flanking sequences useful in the vectors of this invention may be obtained by any of several methods well known in the art. Typically, flanking sequences useful herein – other than the IL-1ra-L gene flanking sequences – will have been previously identified by mapping and/or by restriction endonuclease digestion and  
5 can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full nucleotide sequence of a flanking sequence may be known. Here, the flanking sequence may be synthesized using the methods described herein for nucleic acid synthesis or cloning.

Where all or only a portion of the flanking sequence is known, it may be  
10 obtained using PCR and/or by screening a genomic library with a suitable oligonucleotide and/or flanking sequence fragment from the same or another species. Where the flanking sequence is not known, a fragment of DNA containing a flanking sequence may be isolated from a larger piece of DNA that may contain, for example, a coding sequence or even another gene or genes.  
15 Isolation may be accomplished by restriction endonuclease digestion to produce the proper DNA fragment followed by isolation using agarose gel purification, Qiagen® column chromatography (Chatsworth, CA), or other methods known to the skilled artisan. The selection of suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

20 An origin of replication is typically a part of those prokaryotic expression vectors purchased commercially, and the origin aids in the amplification of the vector in a host cell. Amplification of the vector to a certain copy number can, in some cases, be important for the optimal expression of an IL-1ra-L polypeptide. If the vector of choice does not contain an origin of replication site, one may be  
25 chemically synthesized based on a known sequence, and ligated into the vector. For example, the origin of replication from the plasmid pBR322 (New England Biolabs, Beverly, MA) is suitable for most gram-negative bacteria and various origins (*e.g.*, SV40, polyoma, adenovirus, vesicular stomatitis virus (VSV), or papillomaviruses such as HPV or BPV) are useful for cloning vectors in  
30 mammalian cells. Generally, the origin of replication component is not needed

for mammalian expression vectors (for example, the SV40 origin is often used only because it contains the early promoter).

A transcription termination sequence is typically located 3' of the end of a polypeptide coding region and serves to terminate transcription. Usually, a transcription termination sequence in prokaryotic cells is a G-C rich fragment followed by a poly-T sequence. While the sequence is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described herein.

A selectable marker gene element encodes a protein necessary for the survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, tetracycline, or kanamycin for prokaryotic host cells; (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex media. Preferred selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene. A neomycin resistance gene may also be used for selection in prokaryotic and eukaryotic host cells.

Other selection genes may be used to amplify the gene that will be expressed. Amplification is the process wherein genes that are in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Examples of suitable selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and thymidine kinase. The mammalian cell transformants are placed under selection pressure wherein only the transformants are uniquely adapted to survive by virtue of the selection gene present in the vector. Selection pressure is imposed by culturing the transformed cells under conditions in which the concentration of selection agent in the medium is successively changed, thereby leading to the amplification of both the selection gene and the DNA that encodes an IL-1ra-L polypeptide. As a result, increased quantities of IL-1ra-L polypeptide are synthesized from the amplified DNA.

A ribosome binding site is usually necessary for translation initiation of mRNA and is characterized by a Shine-Dalgarno sequence (prokaryotes) or a Kozak sequence (eukaryotes). The element is typically located 3' to the promoter and 5' to the coding sequence of an IL-1ra-L polypeptide to be expressed. The Shine-Dalgarno sequence is varied but is typically a polypurine (*i.e.*, having a high A-G content). Many Shine-Dalgarno sequences have been identified, each of which can be readily synthesized using methods set forth herein and used in a prokaryotic vector.

A leader, or signal, sequence may be used to direct an IL-1ra-L polypeptide out of the host cell. Typically, a nucleotide sequence encoding the signal sequence is positioned in the coding region of an IL-1ra-L nucleic acid molecule, or directly at the 5' end of an IL-1ra-L polypeptide coding region. Many signal sequences have been identified, and any of those that are functional in the selected host cell may be used in conjunction with an IL-1ra-L nucleic acid molecule. Therefore, a signal sequence may be homologous (naturally occurring) or heterologous to the IL-1ra-L nucleic acid molecule. Additionally, a signal sequence may be chemically synthesized using methods described herein. In most cases, the secretion of an IL-1ra-L polypeptide from the host cell via the presence of a signal peptide will result in the removal of the signal peptide from the secreted IL-1ra-L polypeptide. The signal sequence may be a component of the vector, or it may be a part of an IL-1ra-L nucleic acid molecule that is inserted into the vector.

Included within the scope of this invention is the use of either a nucleotide sequence encoding a native IL-1ra-L polypeptide signal sequence joined to an IL-1ra-L polypeptide coding region or a nucleotide sequence encoding a heterologous signal sequence joined to an IL-1ra-L polypeptide coding region. The heterologous signal sequence selected should be one that is recognized and processed, *i.e.*, cleaved by a signal peptidase, by the host cell. For prokaryotic host cells that do not recognize and process the native IL-1ra-L polypeptide signal sequence, the signal sequence is substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, or

heat-stable enterotoxin II leaders. For yeast secretion, the native IL-1ra-L polypeptide signal sequence may be substituted by the yeast invertase, alpha factor, or acid phosphatase leaders. In mammalian cell expression the native signal sequence is satisfactory, although other mammalian signal sequences may  
5 be suitable.

In some cases, such as where glycosylation is desired in a eukaryotic host cell expression system, one may manipulate the various presequences to improve glycosylation or yield. For example, one may alter the peptidase cleavage site of a particular signal peptide, or add pro-sequences, which also may affect  
10 glycosylation. The final protein product may have, in the -1 position (relative to the first amino acid of the mature protein) one or more additional amino acids incident to expression, which may not have been totally removed. For example, the final protein product may have one or two amino acid residues found in the peptidase cleavage site, attached to the amino-terminus. Alternatively, use of  
15 some enzyme cleavage sites may result in a slightly truncated form of the desired IL-1ra-L polypeptide, if the enzyme cuts at such area within the mature polypeptide.

In many cases, transcription of a nucleic acid molecule is increased by the presence of one or more introns in the vector; this is particularly true where a  
20 polypeptide is produced in eukaryotic host cells, especially mammalian host cells. The introns used may be naturally occurring within the IL-1ra-L gene especially where the gene used is a full-length genomic sequence or a fragment thereof. Where the intron is not naturally occurring within the gene (as for most cDNAs), the intron may be obtained from another source. The position of the intron with  
25 respect to flanking sequences and the IL-1ra-L gene is generally important, as the intron must be transcribed to be effective. Thus, when an IL-1ra-L cDNA molecule is being transcribed, the preferred position for the intron is 3' to the transcription start site and 5' to the poly-A transcription termination sequence. Preferably, the intron or introns will be located on one side or the other (*i.e.*, 5' or  
30 3') of the cDNA such that it does not interrupt the coding sequence. Any intron from any source, including viral, prokaryotic and eukaryotic (plant or animal)

organisms, may be used to practice this invention, provided that it is compatible with the host cell into which it is inserted. Also included herein are synthetic introns. Optionally, more than one intron may be used in the vector.

The expression and cloning vectors of the present invention will typically  
5 contain a promoter that is recognized by the host organism and operably linked to the molecule encoding the IL-1ra-L polypeptide. Promoters are untranscribed sequences located upstream (*i.e.*, 5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription of the structural gene. Promoters are conventionally grouped into one of two classes:  
10 inducible promoters and constitutive promoters. Inducible promoters initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature. Constitutive promoters, on the other hand, initiate continual gene product production; that is, there is little or no control over gene  
15 expression. A large number of promoters, recognized by a variety of potential host cells, are well known. A suitable promoter is operably linked to the DNA encoding IL-1ra-L polypeptide by removing the promoter from the source DNA by restriction enzyme digestion and inserting the desired promoter sequence into the vector. The native IL-1ra-L promoter sequence may be used to direct  
20 amplification and/or expression of an IL-1ra-L nucleic acid molecule. A heterologous promoter is preferred, however, if it permits greater transcription and higher yields of the expressed protein as compared to the native promoter, and if it is compatible with the host cell system that has been selected for use.

Promoters suitable for use with prokaryotic hosts include the beta-lactamase and lactose promoter systems; alkaline phosphatase; a tryptophan (trp) promoter system; and hybrid promoters such as the tac promoter. Other known bacterial promoters are also suitable. Their sequences have been published, thereby enabling one skilled in the art to ligate them to the desired DNA sequence, using linkers or adapters as needed to supply any useful restriction sites.

30 Suitable promoters for use with yeast hosts are also well known in the art. Yeast enhancers are advantageously used with yeast promoters. Suitable

promoters for use with mammalian host cells are well known and include, but are not limited to, those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, retroviruses, hepatitis-B virus and most preferably Simian Virus 40 (SV40). Other suitable mammalian promoters include heterologous mammalian promoters, for example, heat-shock promoters and the actin promoter.

Additional promoters which may be of interest in controlling IL-1ra-L gene expression include, but are not limited to: the SV40 early promoter region (Bernoist and Chambon, 1981, *Nature* 290:304-10); the CMV promoter; the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, *et al.*, 1980, *Cell* 22:787-97); the herpes thymidine kinase promoter (Wagner *et al.*, 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1444-45); the regulatory sequences of the metallothioneine gene (Brinster *et al.*, 1982, *Nature* 296:39-42); prokaryotic expression vectors such as the beta-lactamase promoter (Villa-Kamaroff *et al.*, 1978, *Proc. Natl. Acad. Sci. U.S.A.*, 75:3727-31); or the tac promoter (DeBoer *et al.*, 1983, *Proc. Natl. Acad. Sci. U.S.A.*, 80:21-25). Also of interest are the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: the elastase I gene control region which is active in pancreatic acinar cells (Swift *et al.*, 1984, *Cell* 38:639-46; Ornitz *et al.*, 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409 (1986); MacDonald, 1987, *Hepatology* 7:425-515); the insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-22); the immunoglobulin gene control region which is active in lymphoid cells (Grosschedl *et al.*, 1984, *Cell* 38:647-58; Adames *et al.*, 1985, *Nature* 318:533-38; Alexander *et al.*, 1987, *Mol. Cell. Biol.*, 7:1436-44); the mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder *et al.*, 1986, *Cell* 45:485-95); the albumin gene control region which is active in liver (Pinkert *et al.*, 1987, *Genes and Devel.* 1:268-76); the alpha-feto-protein gene control region which is active in liver (Krumlauf *et al.*, 1985, *Mol. Cell. Biol.*, 5:1639-48; Hammer *et al.*, 1987, *Science* 235:53-58); the alpha 1-

antitrypsin gene control region which is active in the liver (Kelsey *et al.*, 1987, *Genes and Devel.* 1:161-71); the beta-globin gene control region which is active in myeloid cells (Mogram *et al.*, 1985, *Nature* 315:338-40; Kollias *et al.*, 1986, *Cell* 46:89-94); the myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead *et al.*, 1987, *Cell* 48:703-12); the myosin light chain-2 gene control region which is active in skeletal muscle (Sani, 1985, *Nature* 314:283-86); and the gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason *et al.*, 1986, *Science* 234:1372-78).

An enhancer sequence may be inserted into the vector to increase the transcription of a DNA encoding an IL-1ra-L polypeptide of the present invention by higher eukaryotes. Enhancers are cis-acting elements of DNA, usually about 10-300 bp in length, that act on the promoter to increase transcription. Enhancers are relatively orientation and position independent. They have been found 5' and 3' to the transcription unit. Several enhancer sequences available from mammalian genes are known (*e.g.*, globin, elastase, albumin, alpha-feto-protein and insulin). Typically, however, an enhancer from a virus will be used. The SV40 enhancer, the cytomegalovirus early promoter enhancer, the polyoma enhancer, and adenovirus enhancers are exemplary enhancing elements for the activation of eukaryotic promoters. While an enhancer may be spliced into the vector at a position 5' or 3' to an IL-1ra-L nucleic acid molecule, it is typically located at a site 5' from the promoter.

Expression vectors of the invention may be constructed from a starting vector such as a commercially available vector. Such vectors may or may not contain all of the desired flanking sequences. Where one or more of the flanking sequences described herein are not already present in the vector, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the flanking sequences are well known to one skilled in the art.

Preferred vectors for practicing this invention are those which are compatible with bacterial, insect, and mammalian host cells. Such vectors include, *inter alia*, pCRII, pCR3, and pcDNA3.1 (Invitrogen, San Diego, CA),

pBSII (Stratagene, La Jolla, CA), pET15 (Novagen, Madison, WI), pGEX (Pharmacia Biotech, Piscataway, NJ), pEGFP-N2 (Clontech, Palo Alto, CA), pETL (BlueBacII, Invitrogen), pDSR-alpha (PCT Pub. No. WO 90/14363) and pFastBacDual (Gibco-BRL, Grand Island, NY).

5 Additional suitable vectors include, but are not limited to, cosmids, plasmids, or modified viruses, but it will be appreciated that the vector system must be compatible with the selected host cell. Such vectors include, but are not limited to plasmids such as Bluescript<sup>®</sup> plasmid derivatives (a high copy number ColE1-based phagemid, Stratagene Cloning Systems, La Jolla CA), PCR cloning  
10 plasmids designed for cloning Taq-amplified PCR products (*e.g.*, TOPO<sup>™</sup> TA Cloning<sup>®</sup> Kit, PCR2.1<sup>®</sup> plasmid derivatives, Invitrogen, Carlsbad, CA), and mammalian, yeast or virus vectors such as a baculovirus expression system (pBacPAK plasmid derivatives, Clontech, Palo Alto, CA).

After the vector has been constructed and a nucleic acid molecule  
15 encoding an IL-1ra-L polypeptide has been inserted into the proper site of the vector, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression. The transformation of an expression vector for an IL-1ra-L polypeptide into a selected host cell may be accomplished by well known methods including methods such as transfection, infection,  
20 calcium chloride, electroporation, microinjection, lipofection, DEAE-dextran method, or other known techniques. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook *et al.*, *supra*.

25 Host cells may be prokaryotic host cells (such as *E. coli*) or eukaryotic host cells (such as a yeast, insect, or vertebrate cell). The host cell, when cultured under appropriate conditions, synthesizes an IL-1ra-L polypeptide which can subsequently be collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not secreted). The  
30 selection of an appropriate host cell will depend upon various factors, such as desired expression levels, polypeptide modifications that are desirable or



necessary for activity (such as glycosylation or phosphorylation) and ease of folding into a biologically active molecule.

A number of suitable host cells are known in the art and many are available from the American Type Culture Collection (ATCC), Manassas, VA. Examples include, but are not limited to, mammalian cells, such as Chinese hamster ovary cells (CHO), CHO DHFR(-) cells (Urlaub *et al.*, 1980, *Proc. Natl. Acad. Sci. U.S.A.* 97:4216-20), human embryonic kidney (HEK) 293 or 293T cells, or 3T3 cells. The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening, product production, and purification are known in the art. Other suitable mammalian cell lines, are the monkey COS-1 and COS-7 cell lines, and the CV-1 cell line. Further exemplary mammalian host cells include primate cell lines and rodent cell lines, including transformed cell lines. Normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, as well as primary explants, are also suitable. Candidate cells may be genotypically deficient in the selection gene, or may contain a dominantly acting selection gene. Other suitable mammalian cell lines include but are not limited to, mouse neuroblastoma N2A cells, HeLa, mouse L-929 cells, 3T3 lines derived from Swiss, Balb-c or NIH mice, BHK or HaK hamster cell lines. Each of these cell lines is known by and available to those skilled in the art of protein expression.

Similarly useful as host cells suitable for the present invention are bacterial cells. For example, the various strains of *E. coli* (e.g., HB101, DH5 $\alpha$ , DH10, and MC1061) are well-known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Pseudomonas spp.*, other *Bacillus spp.*, *Streptomyces spp.*, and the like may also be employed in this method.

Many strains of yeast cells known to those skilled in the art are also available as host cells for the expression of the polypeptides of the present invention. Preferred yeast cells include, for example, *Saccharomyces cerevisiae* and *Pichia pastoris*.

Additionally, where desired, insect cell systems may be utilized in the methods of the present invention. Such systems are described, for example, in

Kitts *et al.*, 1993, *Biotechniques*, 14:810-17; Lucklow, 1993, *Curr. Opin. Biotechnol.* 4:564-72; and Lucklow *et al.*, 1993, *J. Virol.*, 67:4566-79. Preferred insect cells are Sf-9 and Hi5 (Invitrogen).

One may also use transgenic animals to express glycosylated IL-1ra-L polypeptides. For example, one may use a transgenic milk-producing animal (a cow or goat, for example) and obtain the present glycosylated polypeptide in the animal milk. One may also use plants to produce IL-1ra-L polypeptides, however, in general, the glycosylation occurring in plants is different from that produced in mammalian cells, and may result in a glycosylated product which is not suitable for human therapeutic use.

#### Polypeptide Production

Host cells comprising an IL-1ra-L polypeptide expression vector may be cultured using standard media well known to the skilled artisan. The media will usually contain all nutrients necessary for the growth and survival of the cells. Suitable media for culturing *E. coli* cells include, for example, Luria Broth (LB) and/or Terrific Broth (TB). Suitable media for culturing eukaryotic cells include Roswell Park Memorial Institute medium 1640 (RPMI 1640), Minimal Essential Medium (MEM) and/or Dulbecco's Modified Eagle Medium (DMEM), all of which may be supplemented with serum and/or growth factors as necessary for the particular cell line being cultured. A suitable medium for insect cultures is Grace's medium supplemented with yeastolate, lactalbumin hydrolysate, and/or fetal calf serum as necessary.

Typically, an antibiotic or other compound useful for selective growth of transfected or transformed cells is added as a supplement to the media. The compound to be used will be dictated by the selectable marker element present on the plasmid with which the host cell was transformed. For example, where the selectable marker element is kanamycin resistance, the compound added to the culture medium will be kanamycin. Other compounds for selective growth include ampicillin, tetracycline, and neomycin.

The amount of an IL-1ra-L polypeptide produced by a host cell can be evaluated using standard methods known in the art. Such methods include, without limitation, Western blot analysis, SDS-polyacrylamide gel electrophoresis, non-denaturing gel electrophoresis, High Performance Liquid  
5 Chromatography (HPLC) separation, immunoprecipitation, and/or activity assays such as DNA binding gel shift assays.

If an IL-1ra-L polypeptide has been designed to be secreted from the host cells, the majority of polypeptide may be found in the cell culture medium. If however, the IL-1ra-L polypeptide is not secreted from the host cells, it will be  
10 present in the cytoplasm and/or the nucleus (for eukaryotic host cells) or in the cytosol (for gram-negative bacteria host cells).

For an IL-1ra-L polypeptide situated in the host cell cytoplasm and/or nucleus (for eukaryotic host cells) or in the cytosol (for bacterial host cells), the intracellular material (including inclusion bodies for gram-negative bacteria) can  
15 be extracted from the host cell using any standard technique known to the skilled artisan. For example, the host cells can be lysed to release the contents of the periplasm/cytoplasm by French press, homogenization, and/or sonication followed by centrifugation.

If an IL-1ra-L polypeptide has formed inclusion bodies in the cytosol, the  
20 inclusion bodies can often bind to the inner and/or outer cellular membranes and thus will be found primarily in the pellet material after centrifugation. The pellet material can then be treated at pH extremes or with a chaotropic agent such as a detergent, guanidine, guanidine derivatives, urea, or urea derivatives in the presence of a reducing agent such as dithiothreitol at alkaline pH or tris  
25 carboxyethyl phosphine at acid pH to release, break apart, and solubilize the inclusion bodies. The solubilized IL-1ra-L polypeptide can then be analyzed using gel electrophoresis, immunoprecipitation, or the like. If it is desired to isolate the IL-1ra-L polypeptide, isolation may be accomplished using standard methods such as those described herein and in Marston *et al.*, 1990, *Meth. Enz.*,  
30 182:264-75.

In some cases, an IL-1ra-L polypeptide may not be biologically active upon isolation. Various methods for “refolding” or converting the polypeptide to its tertiary structure and generating disulfide linkages can be used to restore biological activity. Such methods include exposing the solubilized polypeptide to a pH usually above 7 and in the presence of a particular concentration of a chaotrope. The selection of chaotrope is very similar to the choices used for inclusion body solubilization, but usually the chaotrope is used at a lower concentration and is not necessarily the same as chaotropes used for the solubilization. In most cases the refolding/oxidation solution will also contain a reducing agent or the reducing agent plus its oxidized form in a specific ratio to generate a particular redox potential allowing for disulfide shuffling to occur in the formation of the protein’s cysteine bridges. Some of the commonly used redox couples include cysteine/cystamine, glutathione (GSH)/dithiobis GSH, cupric chloride, dithiothreitol(DTT)/dithiane DTT, and 2-2-mercaptoethanol(bME)/dithio-b(ME). In many instances, a cosolvent may be used or may be needed to increase the efficiency of the refolding, and the more common reagents used for this purpose include glycerol, polyethylene glycol of various molecular weights, arginine and the like.

If inclusion bodies are not formed to a significant degree upon expression of an IL-1ra-L polypeptide, then the polypeptide will be found primarily in the supernatant after centrifugation of the cell homogenate. The polypeptide may be further isolated from the supernatant using methods such as those described herein.

The purification of an IL-1ra-L polypeptide from solution can be accomplished using a variety of techniques. If the polypeptide has been synthesized such that it contains a tag such as Hexahistidine (IL-1ra-L polypeptide/hexaHis) or other small peptide such as FLAG (Eastman Kodak Co., New Haven, CT) or *myc* (Invitrogen, Carlsbad, CA) at either its carboxyl- or amino-terminus, it may be purified in a one-step process by passing the solution through an affinity column where the column matrix has a high affinity for the tag.

For example, polyhistidine binds with great affinity and specificity to nickel. Thus, an affinity column of nickel (such as the Qiagen<sup>®</sup> nickel columns) can be used for purification of IL-1ra-L polypeptide/polyHis. *See, e.g., Current Protocols in Molecular Biology* § 10.11.8 (Ausubel *et al.*, eds., Green Publishers Inc. and Wiley and Sons 1993).

Additionally, IL-1RA-L polypeptides may be purified through the use of a monoclonal antibody that is capable of specifically recognizing and binding to an IL-1ra-L polypeptide.

Other suitable procedures for purification include, without limitation, affinity chromatography, immunoaffinity chromatography, ion exchange chromatography, molecular sieve chromatography, HPLC, electrophoresis (including native gel electrophoresis) followed by gel elution, and preparative isoelectric focusing ("Isoprime" machine/technique, Hoefer Scientific, San Francisco, CA). In some cases, two or more purification techniques may be combined to achieve increased purity.

IL-1ra-L polypeptides may also be prepared by chemical synthesis methods (such as solid phase peptide synthesis) using techniques known in the art such as those set forth by Merrifield *et al.*, 1963, *J. Am. Chem. Soc.* 85:2149; Houghten *et al.*, 1985, *Proc Natl Acad. Sci. USA* 82:5132; and Stewart and Young, *Solid Phase Peptide Synthesis* (Pierce Chemical Co. 1984). Such polypeptides may be synthesized with or without a methionine on the amino-terminus. Chemically synthesized IL-1ra-L polypeptides may be oxidized using methods set forth in these references to form disulfide bridges. Chemically synthesized IL-1ra-L polypeptides are expected to have comparable biological activity to the corresponding IL-1ra-L polypeptides produced recombinantly or purified from natural sources, and thus may be used interchangeably with a recombinant or natural IL-1ra-L polypeptide.

Another means of obtaining IL-1ra-L polypeptide is via purification from biological samples such as source tissues and/or fluids in which the IL-1ra-L polypeptide is naturally found. Such purification can be conducted using methods for protein purification as described herein. The presence of the IL-1ra-L

polypeptide during purification may be monitored, for example, using an antibody prepared against recombinantly produced IL-1ra-L polypeptide or peptide fragments thereof.

5 A number of additional methods for producing nucleic acids and polypeptides are known in the art, and the methods can be used to produce polypeptides having specificity for IL-1ra-L polypeptide. *See, e.g.*, Roberts *et al.*, 1997, *Proc. Natl. Acad. Sci. U.S.A.* 94:12297-303, which describes the production of fusion proteins between an mRNA and its encoded peptide. *See also*, Roberts, 1999, *Curr. Opin. Chem. Biol.* 3:268-73. Additionally, U.S. Patent No. 5,824,469  
10 describes methods for obtaining oligonucleotides capable of carrying out a specific biological function. The procedure involves generating a heterogeneous pool of oligonucleotides, each having a 5' randomized sequence, a central preselected sequence, and a 3' randomized sequence. The resulting heterogeneous pool is introduced into a population of cells that do not exhibit the  
15 desired biological function. Subpopulations of the cells are then screened for those that exhibit a predetermined biological function. From that subpopulation, oligonucleotides capable of carrying out the desired biological function are isolated.

U.S. Patent Nos. 5,763,192; 5,814,476; 5,723,323; and 5,817,483 describe  
20 processes for producing peptides or polypeptides. This is done by producing stochastic genes or fragments thereof, and then introducing these genes into host cells which produce one or more proteins encoded by the stochastic genes. The host cells are then screened to identify those clones producing peptides or polypeptides having the desired activity.

25 Another method for producing peptides or polypeptides is described in PCT/US98/20094 (WO99/15650) filed by Athersys, Inc. Known as "Random Activation of Gene Expression for Gene Discovery" (RAGE-GD), the process involves the activation of endogenous gene expression or over-expression of a gene by *in situ* recombination methods. For example, expression of an  
30 endogenous gene is activated or increased by integrating a regulatory sequence into the target cell which is capable of activating expression of the gene by non-

homologous or illegitimate recombination. The target DNA is first subjected to radiation, and a genetic promoter inserted. The promoter eventually locates a break at the front of a gene, initiating transcription of the gene. This results in expression of the desired peptide or polypeptide.

5           It will be appreciated that these methods can also be used to create comprehensive IL-1ra-L polypeptide expression libraries, which can subsequently be used for high throughput phenotypic screening in a variety of assays, such as biochemical assays, cellular assays, and whole organism assays (*e.g.*, plant, mouse, etc.).

10

### Synthesis

It will be appreciated by those skilled in the art that the nucleic acid and polypeptide molecules described herein may be produced by recombinant and other means.

15

### Selective Binding Agents

The term "selective binding agent" refers to a molecule that has specificity for one or more IL-1ra-L polypeptides. Suitable selective binding agents include, but are not limited to, antibodies and derivatives thereof, polypeptides, and small molecules. Suitable selective binding agents may be prepared using methods known in the art. An exemplary IL-1RA-L polypeptide selective binding agent of the present invention is capable of binding a certain portion of the IL-1RA-L polypeptide thereby inhibiting the binding of the polypeptide to an IL-1ra-L polypeptide receptor.

25           Selective binding agents such as antibodies and antibody fragments that bind IL-1ra-L polypeptides are within the scope of the present invention. The antibodies may be polyclonal including monospecific polyclonal; monoclonal (MAbs); recombinant; chimeric; humanized, such as CDR-grafted; human; single chain; and/or bispecific; as well as fragments; variants; or derivatives thereof.

30           Antibody fragments include those portions of the antibody that bind to an epitope on the IL-1RA-L polypeptide. Examples of such fragments include Fab and

F(ab') fragments generated by enzymatic cleavage of full-length antibodies. Other binding fragments include those generated by recombinant DNA techniques, such as the expression of recombinant plasmids containing nucleic acid sequences encoding antibody variable regions.

5 Polyclonal antibodies directed toward an IL-1ra-L polypeptide generally are produced in animals (*e.g.*, rabbits or mice) by means of multiple subcutaneous or intraperitoneal injections of IL-1ra-L polypeptide and an adjuvant. It may be useful to conjugate an IL-1ra-L polypeptide to a carrier protein that is immunogenic in the species to be immunized, such as keyhole limpet  
10 hemocyanin, serum, albumin, bovine thyroglobulin, or soybean trypsin inhibitor. Also, aggregating agents such as alum are used to enhance the immune response. After immunization, the animals are bled and the serum is assayed for anti-IL-1ra-L antibody titer.

Monoclonal antibodies directed toward IL-1ra-L polypeptides are  
15 produced using any method that provides for the production of antibody molecules by continuous cell lines in culture. Examples of suitable methods for preparing monoclonal antibodies include the hybridoma methods of Kohler *et al.*, 1975, *Nature* 256:495-97 and the human B-cell hybridoma method (Kozbor, 1984, *J. Immunol.* 133:3001; Brodeur *et al.*, *Monoclonal Antibody Production*  
20 *Techniques and Applications* 51-63 (Marcel Dekker, Inc., 1987). Also provided by the invention are hybridoma cell lines that produce monoclonal antibodies reactive with IL-1ra-L polypeptides.

Monoclonal antibodies of the invention may be modified for use as  
therapeutics. One embodiment is a "chimeric" antibody in which a portion of the  
25 heavy (H) and/or light (L) chain is identical with or homologous to a corresponding sequence in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is/are identical with or homologous to a corresponding sequence in antibodies derived from another species or belonging to another antibody class or  
30 subclass. Also included are fragments of such antibodies, so long as they exhibit



the desired biological activity. See U.S. Patent No. 4,816,567; Morrison *et al.*, 1985, *Proc. Natl. Acad. Sci.* 81:6851-55.

In another embodiment, a monoclonal antibody of the invention is a "humanized" antibody. Methods for humanizing non-human antibodies are well known in the art. See U.S. Patent Nos. 5,585,089 and 5,693,762. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source that is non-human. Humanization can be performed, for example, using methods described in the art (Jones *et al.*, 1986, *Nature* 321:522-25; Riechmann *et al.*, 1998, *Nature* 332:323-27; Verhoeyen *et al.*, 1988, *Science* 239:1534-36), by substituting at least a portion of a rodent complementarity-determining region (CDR) for the corresponding regions of a human antibody.

Also encompassed by the invention are human antibodies that bind IL-1ra-L polypeptides. Using transgenic animals (*e.g.*, mice) that are capable of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production such antibodies are produced by immunization with an IL-1ra-L polypeptide antigen (*i.e.*, having at least 6 contiguous amino acids), optionally conjugated to a carrier. See, *e.g.*, Jakobovits *et al.*, 1993, *Proc. Natl. Acad. Sci.* 90:2551-55; Jakobovits *et al.*, 1993, *Nature* 362:255-58; Bruggermann *et al.*, 1993, *Year in Immuno.* 7:33. In one method, such transgenic animals are produced by incapacitating the endogenous loci encoding the heavy and light immunoglobulin chains therein, and inserting loci encoding human heavy and light chain proteins into the genome thereof. Partially modified animals, that is those having less than the full complement of modifications, are then cross-bred to obtain an animal having all of the desired immune system modifications. When administered an immunogen, these transgenic animals produce antibodies with human (rather than, *e.g.*, murine) amino acid sequences, including variable regions which are immunospecific for these antigens. See PCT App. Nos. PCT/US96/05928 and PCT/US93/06926. Additional methods are described in U.S. Patent No. 5,545,807, PCT App. Nos. PCT/US91/245 and PCT/GB89/01207, and in European Patent Nos. 546073B1 and 546073A1.

Human antibodies can also be produced by the expression of recombinant DNA in host cells or by expression in hybridoma cells as described herein.

In an alternative embodiment, human antibodies can also be produced from phage-display libraries (Hoogenboom *et al.*, 1991, *J. Mol. Biol.* 227:381; 5 Marks *et al.*, 1991, *J. Mol. Biol.* 222:581). These processes mimic immune selection through the display of antibody repertoires on the surface of filamentous bacteriophage, and subsequent selection of phage by their binding to an antigen of choice. One such technique is described in PCT App. No. PCT/US98/17364, which describes the isolation of high affinity and functional agonistic antibodies 10 for MPL- and msk- receptors using such an approach.

Chimeric, CDR grafted, and humanized antibodies are typically produced by recombinant methods. Nucleic acids encoding the antibodies are introduced into host cells and expressed using materials and procedures described herein. In a preferred embodiment, the antibodies are produced in mammalian host cells, 15 such as CHO cells. Monoclonal (*e.g.*, human) antibodies may be produced by the expression of recombinant DNA in host cells or by expression in hybridoma cells as described herein.

The anti-IL-1ra-L antibodies of the invention may be employed in any known assay method, such as competitive binding assays, direct and indirect 20 sandwich assays, and immunoprecipitation assays (Sola, *Monoclonal Antibodies: A Manual of Techniques* 147-158 (CRC Press, Inc., 1987)) for the detection and quantitation of IL-1ra-L polypeptides. The antibodies will bind IL-1ra-L polypeptides with an affinity that is appropriate for the assay method being employed.

25 For diagnostic applications, in certain embodiments, anti-IL-1ra-L antibodies may be labeled with a detectable moiety. The detectable moiety can be any one that is capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as <sup>3</sup>H, <sup>14</sup>C, <sup>32</sup>P, <sup>35</sup>S, <sup>125</sup>I, <sup>99</sup>Tc, <sup>111</sup>In, or <sup>67</sup>Ga; a fluorescent or chemiluminescent 30 compound, such as fluorescein isothiocyanate, rhodamine, or luciferin; or an

enzyme, such as alkaline phosphatase,  $\beta$ -galactosidase, or horseradish peroxidase (Bayer, *et al.*, 1990, *Meth. Enz.* 184:138-63).

Competitive binding assays rely on the ability of a labeled standard (*e.g.*, an IL-1ra-L polypeptide, or an immunologically reactive portion thereof) to  
5 compete with the test sample analyte (an IL-1ra-L polypeptide) for binding with a limited amount of anti-IL-1ra-L antibody. The amount of an IL-1ra-L polypeptide in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of  
10 standard that becomes bound, the antibodies typically are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the standard and analyte which remain unbound.

Sandwich assays typically involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be  
15 detected and/or quantitated. In a sandwich assay, the test sample analyte is typically bound by a first antibody which is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three-part complex. *See, e.g.*, U.S. Patent No. 4,376,110. The second antibody may  
20 itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assays). For example, one type of sandwich assay is an enzyme-linked immunosorbent assay (ELISA), in which case the detectable moiety is an enzyme.

The selective binding agents, including anti-IL-1ra-L antibodies, are also  
25 useful for *in vivo* imaging. An antibody labeled with a detectable moiety may be administered to an animal, preferably into the bloodstream, and the presence and location of the labeled antibody in the host assayed. The antibody may be labeled with any moiety that is detectable in an animal, whether by nuclear magnetic resonance, radiology, or other detection means known in the art.

30 Selective binding agents of the invention, including antibodies, may be used as therapeutics. These therapeutic agents are generally agonists or

antagonists, in that they either enhance or reduce, respectively, at least one of the biological activities of an IL-1ra-L polypeptide. In one embodiment, antagonist antibodies of the invention are antibodies or binding fragments thereof which are capable of specifically binding to an IL-1ra-L polypeptide and which are capable of inhibiting or eliminating the functional activity of an IL-1ra-L polypeptide *in vivo* or *in vitro*. In preferred embodiments, the selective binding agent, *e.g.*, an antagonist antibody, will inhibit the functional activity of an IL-1ra-L polypeptide by at least about 50%, and preferably by at least about 80%. In another embodiment, the selective binding agent may be an anti-IL-1ra-L polypeptide antibody that is capable of interacting with an IL-1ra-L polypeptide binding partner (a ligand or receptor) thereby inhibiting or eliminating IL-1ra-L polypeptide activity *in vitro* or *in vivo*. Selective binding agents, including agonist and antagonist anti-IL-1ra-L polypeptide antibodies, are identified by screening assays that are well known in the art.

The invention also relates to a kit comprising IL-1ra-L selective binding agents (such as antibodies) and other reagents useful for detecting IL-1ra-L polypeptide levels in biological samples. Such reagents may include a detectable label, blocking serum, positive and negative control samples, and detection reagents.

20

#### Microarrays

It will be appreciated that DNA microarray technology can be utilized in accordance with the present invention. DNA microarrays are miniature, high-density arrays of nucleic acids positioned on a solid support, such as glass. Each cell or element within the array contains numerous copies of a single nucleic acid species that acts as a target for hybridization with a complementary nucleic acid sequence (*e.g.*, mRNA). In expression profiling using DNA microarray technology, mRNA is first extracted from a cell or tissue sample and then converted enzymatically to fluorescently labeled cDNA. This material is hybridized to the microarray and unbound cDNA is removed by washing. The expression of discrete genes represented on the array is then visualized by

quantitating the amount of labeled cDNA that is specifically bound to each target nucleic acid molecule. In this way, the expression of thousands of genes can be quantitated in a high throughput, parallel manner from a single sample of biological material.

5           This high throughput expression profiling has a broad range of applications with respect to the IL-1ra-L molecules of the invention, including, but not limited to: the identification and validation of IL-1ra-L disease-related genes as targets for therapeutics; molecular toxicology of related IL-1ra-L molecules and inhibitors thereof; stratification of populations and generation of  
10 surrogate markers for clinical trials; and enhancing related IL-1ra-L polypeptide small molecule drug discovery by aiding in the identification of selective compounds in high throughput screens.

#### Chemical Derivatives

15           Chemically modified derivatives of IL-1ra-L polypeptides may be prepared by one skilled in the art, given the disclosures described herein. IL-1ra-L polypeptide derivatives are modified in a manner that is different – either in the type or location of the molecules naturally attached to the polypeptide. Derivatives may include molecules formed by the deletion of one or more  
20 naturally-attached chemical groups. The polypeptide comprising the amino acid sequence of SEQ ID NO: 2, or other IL-1ra-L polypeptide, may be modified by the covalent attachment of one or more polymers. For example, the polymer selected is typically water-soluble so that the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment.  
25 Included within the scope of suitable polymers is a mixture of polymers. Preferably, for therapeutic use of the end-product preparation, the polymer will be pharmaceutically acceptable.

          The polymers each may be of any molecular weight and may be branched or unbranched. The polymers each typically have an average molecular weight of  
30 between about 2 kDa to about 100 kDa (the term “about” indicating that in preparations of a water-soluble polymer, some molecules will weigh more, some

less, than the stated molecular weight). The average molecular weight of each polymer is preferably between about 5 kDa and about 50 kDa, more preferably between about 12 kDa and about 40 kDa and most preferably between about 20 kDa and about 35 kDa.

5           Suitable water-soluble polymers or mixtures thereof include, but are not limited to, N-linked or O-linked carbohydrates, sugars, phosphates, polyethylene glycol (PEG) (including the forms of PEG that have been used to derivatize proteins, including mono-(C<sub>1</sub>-C<sub>10</sub>), alkoxy-, or aryloxy-polyethylene glycol), monomethoxy-polyethylene glycol, dextran (such as low molecular weight  
10 dextran of, for example, about 6 kD), cellulose, or other carbohydrate based polymers, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols (*e.g.*, glycerol), and polyvinyl alcohol. Also encompassed by the present invention are bifunctional crosslinking molecules  
15 which may be used to prepare covalently attached IL-1ra-L polypeptide multimers.

          In general, chemical derivatization may be performed under any suitable condition used to react a protein with an activated polymer molecule. Methods for preparing chemical derivatives of polypeptides will generally comprise the  
20 steps of: (a) reacting the polypeptide with the activated polymer molecule (such as a reactive ester or aldehyde derivative of the polymer molecule) under conditions whereby the polypeptide comprising the amino acid sequence of SEQ ID NO: 2, or other IL-1ra-L polypeptide, becomes attached to one or more polymer molecules, and (b) obtaining the reaction products. The optimal reaction  
25 conditions will be determined based on known parameters and the desired result. For example, the larger the ratio of polymer molecules to protein, the greater the percentage of attached polymer molecule. In one embodiment, the IL-1ra-L polypeptide derivative may have a single polymer molecule moiety at the amino-terminus. *See, e.g.*, U.S. Patent No. 5,234,784.

30           The pegylation of a polypeptide may be specifically carried out using any of the pegylation reactions known in the art. Such reactions are described, for

example, in the following references: Francis *et al.*, 1992, *Focus on Growth Factors* 3:4-10; European Patent Nos. 0154316 and 0401384; and U.S. Patent No. 4,179,337. For example, pegylation may be carried out via an acylation reaction or an alkylation reaction with a reactive polyethylene glycol molecule (or an analogous reactive water-soluble polymer) as described herein. For the acylation reactions, a selected polymer should have a single reactive ester group. For reductive alkylation, a selected polymer should have a single reactive aldehyde group. A reactive aldehyde is, for example, polyethylene glycol propionaldehyde, which is water stable, or mono C<sub>1</sub>-C<sub>10</sub> alkoxy or aryloxy derivatives thereof (see U.S. Patent No. 5,252,714).

In another embodiment, IL-1ra-L polypeptides may be chemically coupled to biotin. The biotin/IL-1ra-L polypeptide molecules are then allowed to bind to avidin, resulting in tetravalent avidin/biotin/IL-1ra-L polypeptide molecules. IL-1ra-L polypeptides may also be covalently coupled to dinitrophenol (DNP) or trinitrophenol (TNP) and the resulting conjugates precipitated with anti-DNP or anti-TNP-IgM to form decameric conjugates with a valency of 10.

Generally, conditions that may be alleviated or modulated by the administration of the present IL-1ra-L polypeptide derivatives include those described herein for IL-1ra-L polypeptides. However, the IL-1ra-L polypeptide derivatives disclosed herein may have additional activities, enhanced or reduced biological activity, or other characteristics, such as increased or decreased half-life, as compared to the non-derivatized molecules.

#### Genetically Engineered Non-Human Animals

Additionally included within the scope of the present invention are non-human animals such as mice, rats, or other rodents; rabbits, goats, sheep, or other farm animals, in which the genes encoding native IL-1ra-L polypeptide have been disrupted (*i.e.*, “knocked out”) such that the level of expression of IL-1ra-L polypeptide is significantly decreased or completely abolished. Such animals may be prepared using techniques and methods such as those described in U.S. Patent No. 5,557,032.

The present invention further includes non-human animals such as mice, rats, or other rodents; rabbits, goats, sheep, or other farm animals, in which either the native form of an IL-1ra-L gene for that animal or a heterologous IL-1ra-L gene is over-expressed by the animal, thereby creating a “transgenic” animal.

5 Such transgenic animals may be prepared using well known methods such as those described in U.S. Patent No 5,489,743 and PCT Pub. No. WO 94/28122.

The present invention further includes non-human animals in which the promoter for one or more of the IL-1ra-L polypeptides of the present invention is either activated or inactivated (*e.g.*, by using homologous recombination methods) to alter the level of expression of one or more of the native IL-1ra-L polypeptides.

These non-human animals may be used for drug candidate screening. In such screening, the impact of a drug candidate on the animal may be measured. For example, drug candidates may decrease or increase the expression of the IL-1ra-L gene. In certain embodiments, the amount of IL-1ra-L polypeptide that is produced may be measured after the exposure of the animal to the drug candidate. Additionally, in certain embodiments, one may detect the actual impact of the drug candidate on the animal. For example, over-expression of a particular gene may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate’s ability to decrease expression of the gene or its ability to prevent or inhibit a pathological condition. In other examples, the production of a particular metabolic product such as a fragment of a polypeptide, may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate’s ability to decrease the production of such a metabolic product or its ability to prevent or inhibit a pathological condition.

25

#### Assaying for Other Modulators of IL-1ra-L Polypeptide Activity

In some situations, it may be desirable to identify molecules that are modulators, *i.e.*, agonists or antagonists, of the activity of IL-1ra-L polypeptide. Natural or synthetic molecules that modulate IL-1ra-L polypeptide may be identified using one or more screening assays, such as those described herein.

30



Such molecules may be administered either in an *ex vivo* manner or in an *in vivo* manner by injection, or by oral delivery, implantation device, or the like.

“Test molecule” refers to a molecule that is under evaluation for the ability to modulate (*i.e.*, increase or decrease) the activity of an IL-1ra-L polypeptide.

5 Most commonly, a test molecule will interact directly with an IL-1ra-L polypeptide. However, it is also contemplated that a test molecule may also modulate IL-1ra-L polypeptide activity indirectly, such as by affecting IL-1ra-L gene expression, or by binding to an IL-1ra-L polypeptide binding partner (*e.g.*, receptor or ligand). In one embodiment, a test molecule will bind to an IL-1ra-L  
10 polypeptide with an affinity constant of at least about  $10^{-6}$  M, preferably about  $10^{-8}$  M, more preferably about  $10^{-9}$  M, and even more preferably about  $10^{-10}$  M.

Methods for identifying compounds that interact with IL-1ra-L polypeptides are encompassed by the present invention. In certain embodiments, an IL-1ra-L polypeptide is incubated with a test molecule under conditions that  
15 permit the interaction of the test molecule with an IL-1ra-L polypeptide, and the extent of the interaction is measured. The test molecule can be screened in a substantially purified form or in a crude mixture.

In certain embodiments, an IL-1ra-L polypeptide agonist or antagonist may be a protein, peptide, carbohydrate, lipid, or small molecular weight  
20 molecule that interacts with IL-1ra-L polypeptide to regulate its activity. Molecules which regulate IL-1ra-L polypeptide expression include nucleic acids which are complementary to nucleic acids encoding an IL-1ra-L polypeptide, or are complementary to nucleic acids sequences which direct or control the expression of IL-1ra-L polypeptide, and which act as anti-sense regulators of  
25 expression.

Once a test molecule has been identified as interacting with an IL-1ra-L polypeptide, the molecule may be further evaluated for its ability to increase or decrease IL-1ra-L polypeptide activity. The measurement of the interaction of a test molecule with IL-1ra-L polypeptide may be carried out in several formats,  
30 including cell-based binding assays, membrane binding assays, solution-phase assays, and immunoassays. In general, a test molecule is incubated with an IL-

1ra-L polypeptide for a specified period of time, and IL-1ra-L polypeptide activity is determined by one or more assays for measuring biological activity.

The interaction of test molecules with IL-1ra-L polypeptides may also be assayed directly using polyclonal or monoclonal antibodies in an immunoassay.

- 5 Alternatively, modified forms of IL-1ra-L polypeptides containing epitope tags as described herein may be used in solution and immunoassays.

In the event that IL-1ra-L polypeptides display biological activity through an interaction with a binding partner (*e.g.*, a receptor or a ligand), a variety of *in vitro* assays may be used to measure the binding of an IL-1ra-L polypeptide to the  
10 corresponding binding partner (such as a selective binding agent, receptor, or ligand). These assays may be used to screen test molecules for their ability to increase or decrease the rate and/or the extent of binding of an IL-1ra-L polypeptide to its binding partner. In one assay, an IL-1ra-L polypeptide is immobilized in the wells of a microtiter plate. Radiolabeled IL-1ra-L polypeptide  
15 binding partner (for example, iodinated IL-1ra-L polypeptide binding partner) and a test molecule can then be added either one at a time (in either order) or simultaneously to the wells. After incubation, the wells can be washed and counted for radioactivity, using a scintillation counter, to determine the extent to which the binding partner bound to the IL-1ra-L polypeptide. Typically, a  
20 molecule will be tested over a range of concentrations, and a series of control wells lacking one or more elements of the test assays can be used for accuracy in the evaluation of the results. An alternative to this method involves reversing the “positions” of the proteins, *i.e.*, immobilizing IL-1ra-L polypeptide binding partner to the microtiter plate wells, incubating with the test molecule and  
25 radiolabeled IL-1ra-L polypeptide, and determining the extent of IL-1ra-L polypeptide binding. *See, e.g., Current Protocols in Molecular Biology*, chap. 18 (Ausubel *et al.*, eds., Green Publishers Inc. and Wiley and Sons 1995).

As an alternative to radiolabeling, an IL-1ra-L polypeptide or its binding partner may be conjugated to biotin, and the presence of biotinylated protein can  
30 then be detected using streptavidin linked to an enzyme, such as horse radish peroxidase (HRP) or alkaline phosphatase (AP), which can be detected

colorometrically, or by fluorescent tagging of streptavidin. An antibody directed to an IL-1ra-L polypeptide or to an IL-1ra-L polypeptide binding partner, and which is conjugated to biotin, may also be used for purposes of detection following incubation of the complex with enzyme-linked streptavidin linked to AP or HRP.

A IL-1ra-L polypeptide or an IL-1ra-L polypeptide binding partner can also be immobilized by attachment to agarose beads, acrylic beads, or other types of such inert solid phase substrates. The substrate-protein complex can be placed in a solution containing the complementary protein and the test compound. After incubation, the beads can be precipitated by centrifugation, and the amount of binding between an IL-1ra-L polypeptide and its binding partner can be assessed using the methods described herein. Alternatively, the substrate-protein complex can be immobilized in a column with the test molecule and complementary protein passing through the column. The formation of a complex between an IL-1ra-L polypeptide and its binding partner can then be assessed using any of the techniques described herein (*e.g.*, radiolabelling or antibody binding).

Another *in vitro* assay that is useful for identifying a test molecule which increases or decreases the formation of a complex between an IL-1ra-L polypeptide binding protein and an IL-1ra-L polypeptide binding partner is a surface plasmon resonance detector system such as the BIAcore assay system (Pharmacia, Piscataway, NJ). The BIAcore system is utilized as specified by the manufacturer. This assay essentially involves the covalent binding of either IL-1ra-L polypeptide or an IL-1ra-L polypeptide binding partner to a dextran-coated sensor chip that is located in a detector. The test compound and the other complementary protein can then be injected, either simultaneously or sequentially, into the chamber containing the sensor chip. The amount of complementary protein that binds can be assessed based on the change in molecular mass that is physically associated with the dextran-coated side of the sensor chip, with the change in molecular mass being measured by the detector system.

In some cases, it may be desirable to evaluate two or more test compounds together for their ability to increase or decrease the formation of a complex

between an IL-1ra-L polypeptide and an IL-1ra-L polypeptide binding partner. In these cases, the assays set forth herein can be readily modified by adding such additional test compound(s) either simultaneously with, or subsequent to, the first test compound. The remainder of the steps in the assay are as set forth herein.

5           *In vitro* assays such as those described herein may be used advantageously to screen large numbers of compounds for an effect on the formation of a complex between an IL-1ra-L polypeptide and IL-1ra-L polypeptide binding partner. The assays may be automated to screen compounds generated in phage display, synthetic peptide, and chemical synthesis libraries.

10           Compounds which increase or decrease the formation of a complex between an IL-1ra-L polypeptide and an IL-1ra-L polypeptide binding partner may also be screened in cell culture using cells and cell lines expressing either IL-1ra-L polypeptide or IL-1ra-L polypeptide binding partner. Cells and cell lines may be obtained from any mammal, but preferably will be from human or other  
15           primate, canine, or rodent sources. The binding of an IL-1ra-L polypeptide to cells expressing IL-1ra-L polypeptide binding partner at the surface is evaluated in the presence or absence of test molecules, and the extent of binding may be determined by, for example, flow cytometry using a biotinylated antibody to an IL-1ra-L polypeptide binding partner. Cell culture assays can be used  
20           advantageously to further evaluate compounds that score positive in protein binding assays described herein.

Cell cultures can also be used to screen the impact of a drug candidate. For example, drug candidates may decrease or increase the expression of the IL-1ra-L gene. In certain embodiments, the amount of IL-1ra-L polypeptide or an  
25           IL-1ra-L polypeptide fragment that is produced may be measured after exposure of the cell culture to the drug candidate. In certain embodiments, one may detect the actual impact of the drug candidate on the cell culture. For example, the over-expression of a particular gene may have a particular impact on the cell culture. In such cases, one may test a drug candidate's ability to increase or decrease the  
30           expression of the gene or its ability to prevent or inhibit a particular impact on the cell culture. In other examples, the production of a particular metabolic product

such as a fragment of a polypeptide, may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease the production of such a metabolic product in a cell culture.

5    Internalizing Proteins

          The *tat* protein sequence (from HIV) can be used to internalize proteins into a cell. *See, e.g., Falwell et al., 1994, Proc. Natl. Acad. Sci. U.S.A. 91:664-68.* For example, an 11 amino acid sequence (Y-G-R-K-K-R-R-Q-R-R-R; SEQ ID NO: 7) of the HIV *tat* protein (termed the "protein transduction domain," or TAT  
10   PDT) has been described as mediating delivery across the cytoplasmic membrane and the nuclear membrane of a cell. *See Schwarze et al., 1999, Science 285:1569-72; and Nagahara et al., 1998, Nat. Med. 4:1449-52.* In these procedures, FITC-constructs (FITC-labeled G-G-G-G-Y-G-R-K-K-R-R-Q-R-R-R; SEQ ID NO: 8), which penetrate tissues following intraperitoneal administration, are prepared, and  
15   the binding of such constructs to cells is detected by fluorescence-activated cell sorting (FACS) analysis. Cells treated with a *tat*- $\beta$ -gal fusion protein will demonstrate  $\beta$ -gal activity. Following injection, expression of such a construct can be detected in a number of tissues, including liver, kidney, lung, heart, and brain tissue. It is believed that such constructs undergo some degree of unfolding  
20   in order to enter the cell, and as such, may require a refolding following entry into the cell.

          It will thus be appreciated that the *tat* protein sequence may be used to internalize a desired polypeptide into a cell. For example, using the *tat* protein sequence, an IL-1ra-L antagonist (such as an anti-IL-1ra-L selective binding  
25   agent, small molecule, soluble receptor, or antisense oligonucleotide) can be administered intracellularly to inhibit the activity of an IL-1ra-L molecule. As used herein, the term "IL-1ra-L molecule" refers to both IL-1ra-L nucleic acid molecules and IL-1ra-L polypeptides as defined herein. Where desired, the IL-1ra-L protein itself may also be internally administered to a cell using these  
30   procedures. *See also, Straus, 1999, Science 285:1466-67.*

#### Cell Source Identification Using IL-1ra-L Polypeptide

In accordance with certain embodiments of the invention, it may be useful to be able to determine the source of a certain cell type associated with an IL-1ra-L polypeptide. For example, it may be useful to determine the origin of a disease or pathological condition as an aid in selecting an appropriate therapy. In certain  
5       embodiments, nucleic acids encoding an IL-1ra-L polypeptide can be used as a probe to identify cells described herein by screening the nucleic acids of the cells with such a probe. In other embodiments, one may use anti-IL-1ra-L polypeptide antibodies to test for the presence of IL-1ra-L polypeptide in cells, and thus,  
10       determine if such cells are of the types described herein.

#### IL-1ra-L Polypeptide Compositions and Administration

Therapeutic compositions are within the scope of the present invention. Such IL-1RA-L polypeptide pharmaceutical compositions may comprise a  
15       therapeutically effective amount of an IL-1ra-L polypeptide or an IL-1ra-L nucleic acid molecule in admixture with a pharmaceutically or physiologically acceptable formulation agent selected for suitability with the mode of administration. Pharmaceutical compositions may comprise a therapeutically effective amount of one or more IL-1ra-L polypeptide selective binding agents in  
20       admixture with a pharmaceutically or physiologically acceptable formulation agent selected for suitability with the mode of administration.

Acceptable formulation materials preferably are nontoxic to recipients at the dosages and concentrations employed.

The pharmaceutical composition may contain formulation materials for  
25       modifying, maintaining, or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption, or penetration of the composition. Suitable formulation materials include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine, or lysine), antimicrobials, antioxidants (such as ascorbic  
30       acid, sodium sulfite, or sodium hydrogen-sulfite), buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates, or other organic acids), bulking agents

(such as mannitol or glycine), chelating agents (such as ethylenediamine tetraacetic acid (EDTA)), complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin, or hydroxypropyl-beta-cyclodextrin), fillers, monosaccharides, disaccharides, and other carbohydrates (such as glucose, mannose, or dextrans), proteins (such as serum albumin, gelatin, or immunoglobulins), coloring, flavoring and diluting agents, emulsifying agents, hydrophilic polymers (such as polyvinylpyrrolidone), low molecular weight polypeptides, salt-forming counterions (such as sodium), preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid, or hydrogen peroxide), solvents (such as glycerin, propylene glycol, or polyethylene glycol), sugar alcohols (such as mannitol or sorbitol), suspending agents, surfactants or wetting agents (such as pluronics; PEG; sorbitan esters; polysorbates such as polysorbate 20 or polysorbate 80; triton; tromethamine; lecithin; cholesterol or tyloxapal), stability enhancing agents (such as sucrose or sorbitol), tonicity enhancing agents (such as alkali metal halides – preferably sodium or potassium chloride – or mannitol sorbitol), delivery vehicles, diluents, excipients and/or pharmaceutical adjuvants. *See Remington's Pharmaceutical Sciences* (18th Ed., A.R. Gennaro, ed., Mack Publishing Company 1990.

The optimal pharmaceutical composition will be determined by a skilled artisan depending upon, for example, the intended route of administration, delivery format, and desired dosage. *See, e.g., Remington's Pharmaceutical Sciences, supra.* Such compositions may influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the IL-1ra-L molecule.

The primary vehicle or carrier in a pharmaceutical composition may be either aqueous or non-aqueous in nature. For example, a suitable vehicle or carrier for injection may be water, physiological saline solution, or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. Other exemplary pharmaceutical compositions comprise Tris buffer of about pH 7.0-8.5, or acetate

buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute. In one embodiment of the present invention, IL-1ra-L polypeptide compositions may be prepared for storage by mixing the selected composition having the desired degree of purity with optional formulation agents (*Remington's*  
5 *Pharmaceutical Sciences, supra*) in the form of a lyophilized cake or an aqueous solution. Further, the IL-1ra-L polypeptide product may be formulated as a lyophilizate using appropriate excipients such as sucrose.

The IL-1ra-L polypeptide pharmaceutical compositions can be selected for parenteral delivery. Alternatively, the compositions may be selected for  
10 inhalation or for delivery through the digestive tract, such as orally. The preparation of such pharmaceutically acceptable compositions is within the skill of the art.

The formulation components are present in concentrations that are acceptable to the site of administration. For example, buffers are used to maintain  
15 the composition at physiological pH or at a slightly lower pH, typically within a pH range of from about 5 to about 8.

When parenteral administration is contemplated, the therapeutic compositions for use in this invention may be in the form of a pyrogen-free, parenterally acceptable, aqueous solution comprising the desired IL-1ra-L  
20 molecule in a pharmaceutically acceptable vehicle. A particularly suitable vehicle for parenteral injection is sterile distilled water in which an IL-1ra-L molecule is formulated as a sterile, isotonic solution, properly preserved. Yet another preparation can involve the formulation of the desired molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds  
25 (such as polylactic acid or polyglycolic acid), beads, or liposomes, that provides for the controlled or sustained release of the product which may then be delivered via a depot injection. Hyaluronic acid may also be used, and this may have the effect of promoting sustained duration in the circulation. Other suitable means for the introduction of the desired molecule include implantable drug delivery  
30 devices.



In one embodiment, a pharmaceutical composition may be formulated for inhalation. For example, IL-1ra-L polypeptide may be formulated as a dry powder for inhalation. IL-1ra-L polypeptide or nucleic acid molecule inhalation solutions may also be formulated with a propellant for aerosol delivery. In yet  
5 another embodiment, solutions may be nebulized. Pulmonary administration is further described in PCT Pub. No. WO 94/20069, which describes the pulmonary delivery of chemically modified proteins.

It is also contemplated that certain formulations may be administered orally. In one embodiment of the present invention, IL-1ra-L polypeptides that  
10 are administered in this fashion can be formulated with or without those carriers customarily used in the compounding of solid dosage forms such as tablets and capsules. For example, a capsule may be designed to release the active portion of the formulation at the point in the gastrointestinal tract when bioavailability is maximized and pre-systemic degradation is minimized. Additional agents can be  
15 included to facilitate absorption of the IL-1ra-L polypeptide. Diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be employed.

Another pharmaceutical composition may involve an effective quantity of IL-1ra-L polypeptides in a mixture with non-toxic excipients that are suitable for  
20 the manufacture of tablets. By dissolving the tablets in sterile water, or another appropriate vehicle, solutions can be prepared in unit-dose form. Suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium carbonate or bicarbonate, lactose, or calcium phosphate; or binding agents, such as starch, gelatin, or acacia; or lubricating agents such as  
25 magnesium stearate, stearic acid, or talc.

Additional IL-1ra-L polypeptide pharmaceutical compositions will be evident to those skilled in the art, including formulations involving IL-1ra-L polypeptides in sustained- or controlled-delivery formulations. Techniques for formulating a variety of other sustained- or controlled-delivery means, such as  
30 liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. *See, e.g.*, PCT/US93/00829,

which describes the controlled release of porous polymeric microparticles for the delivery of pharmaceutical compositions.

Additional examples of sustained-release preparations include semipermeable polymer matrices in the form of shaped articles, *e.g.* films, or  
5 microcapsules. Sustained release matrices may include polyesters, hydrogels, polylactides (U.S. Patent No. 3,773,919 and European Patent No. 058481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman *et al.*, 1983, *Biopolymers* 22:547-56), poly(2-hydroxyethyl-methacrylate) (Langer *et al.*, 1981, *J. Biomed. Mater. Res.* 15:167-277 and Langer, 1982, *Chem. Tech.* 12:98-  
10 105), ethylene vinyl acetate (Langer *et al.*, *supra*) or poly-D(-)-3-hydroxybutyric acid (European Patent No. 133988). Sustained-release compositions may also include liposomes, which can be prepared by any of several methods known in the art. *See, e.g.*, Eppstein *et al.*, 1985, *Proc. Natl. Acad. Sci. USA* 82:3688-92; and European Patent Nos. 036676, 088046, and 143949.

15 The IL-1ra-L pharmaceutical composition to be used for *in vivo* administration typically must be sterile. This may be accomplished by filtration through sterile filtration membranes. Where the composition is lyophilized, sterilization using this method may be conducted either prior to, or following, lyophilization and reconstitution. The composition for parenteral administration  
20 may be stored in lyophilized form or in a solution. In addition, parenteral compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Once the pharmaceutical composition has been formulated, it may be  
25 stored in sterile vials as a solution, suspension, gel, emulsion, solid, or as a dehydrated or lyophilized powder. Such formulations may be stored either in a ready-to-use form or in a form (*e.g.*, lyophilized) requiring reconstitution prior to administration.

In a specific embodiment, the present invention is directed to kits for  
30 producing a single-dose administration unit. The kits may each contain both a first container having a dried protein and a second container having an aqueous

formulation. Also included within the scope of this invention are kits containing single and multi-chambered pre-filled syringes (*e.g.*, liquid syringes and lyosyringes).

The effective amount of an IL-1ra-L pharmaceutical composition to be employed therapeutically will depend, for example, upon the therapeutic context and objectives. One skilled in the art will appreciate that the appropriate dosage levels for treatment will thus vary depending, in part, upon the molecule delivered, the indication for which the IL-1ra-L molecule is being used, the route of administration, and the size (body weight, body surface, or organ size) and condition (the age and general health) of the patient. Accordingly, the clinician may titer the dosage and modify the route of administration to obtain the optimal therapeutic effect. A typical dosage may range from about 0.1  $\mu\text{g/kg}$  to up to about 100 mg/kg or more, depending on the factors mentioned above. In other embodiments, the dosage may range from 0.1  $\mu\text{g/kg}$  up to about 100 mg/kg; or 1  $\mu\text{g/kg}$  up to about 100 mg/kg; or 5  $\mu\text{g/kg}$  up to about 100 mg/kg.

The frequency of dosing will depend upon the pharmacokinetic parameters of the IL-1ra-L molecule in the formulation being used. Typically, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, as two or more doses (which may or may not contain the same amount of the desired molecule) over time, or as a continuous infusion via an implantation device or catheter. Further refinement of the appropriate dosage is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed by them. Appropriate dosages may be ascertained through use of appropriate dose-response data.

The route of administration of the pharmaceutical composition is in accord with known methods, *e.g.*, orally; through injection by intravenous, intraperitoneal, intracerebral (intraparenchymal), intracerebroventricular, intramuscular, intraocular, intraarterial, intraportal, or intralesional routes; by sustained release systems; or by implantation devices. Where desired, the

compositions may be administered by bolus injection or continuously by infusion, or by implantation device.

Alternatively or additionally, the composition may be administered locally via implantation of a membrane, sponge, or other appropriate material onto which  
5 the desired molecule has been absorbed or encapsulated. Where an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of the desired molecule may be via diffusion, timed-release bolus, or continuous administration.

In some cases, it may be desirable to use IL-1ra-L polypeptide  
10 pharmaceutical compositions in an *ex vivo* manner. In such instances, cells, tissues, or organs that have been removed from the patient are exposed to IL-1ra-L polypeptide pharmaceutical compositions after which the cells, tissues, or organs are subsequently implanted back into the patient.

In other cases, an IL-1ra-L polypeptide can be delivered by implanting  
15 certain cells that have been genetically engineered, using methods such as those described herein, to express and secrete the IL-1ra-L polypeptide. Such cells may be animal or human cells, and may be autologous, heterologous, or xenogeneic. Optionally, the cells may be immortalized. In order to decrease the chance of an immunological response, the cells may be encapsulated to avoid infiltration of  
20 surrounding tissues. The encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow the release of the protein product(s) but prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

As discussed herein, it may be desirable to treat isolated cell populations  
25 (such as stem cells, lymphocytes, red blood cells, chondrocytes, neurons, and the like) with one or more IL-1ra-L polypeptides. This can be accomplished by exposing the isolated cells to the polypeptide directly, where it is in a form that is permeable to the cell membrane.

Additional embodiments of the present invention relate to cells and  
30 methods (*e.g.*, homologous recombination and/or other recombinant production methods) for both the *in vitro* production of therapeutic polypeptides and for the

production and delivery of therapeutic polypeptides by gene therapy or cell therapy. Homologous and other recombination methods may be used to modify a cell that contains a normally transcriptionally-silent IL-1ra-L gene, or an under-expressed gene, and thereby produce a cell which expresses therapeutically efficacious amounts of IL-1ra-L polypeptides.

Homologous recombination is a technique originally developed for targeting genes to induce or correct mutations in transcriptionally active genes. Kucherlapati, 1989, *Prog. in Nucl. Acid Res. & Mol. Biol.* 36:301. The basic technique was developed as a method for introducing specific mutations into specific regions of the mammalian genome (Thomas *et al.*, 1986, *Cell* 44:419-28; Thomas and Capecchi, 1987, *Cell* 51:503-12; Doetschman *et al.*, 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:8583-87) or to correct specific mutations within defective genes (Doetschman *et al.*, 1987, *Nature* 330:576-78). Exemplary homologous recombination techniques are described in U.S. Patent No. 5,272,071; European Patent Nos. 9193051 and 505500; PCT/US90/07642, and PCT Pub No. WO 91/09955).

Through homologous recombination, the DNA sequence to be inserted into the genome can be directed to a specific region of the gene of interest by attaching it to targeting DNA. The targeting DNA is a nucleotide sequence that is complementary (homologous) to a region of the genomic DNA. Small pieces of targeting DNA that are complementary to a specific region of the genome are put in contact with the parental strand during the DNA replication process. It is a general property of DNA that has been inserted into a cell to hybridize, and therefore, recombine with other pieces of endogenous DNA through shared homologous regions. If this complementary strand is attached to an oligonucleotide that contains a mutation or a different sequence or an additional nucleotide, it too is incorporated into the newly synthesized strand as a result of the recombination. As a result of the proofreading function, it is possible for the new sequence of DNA to serve as the template. Thus, the transferred DNA is incorporated into the genome.

Attached to these pieces of targeting DNA are regions of DNA that may interact with or control the expression of an IL-1ra-L polypeptide, *e.g.*, flanking sequences. For example, a promoter/enhancer element, a suppressor, or an exogenous transcription modulatory element is inserted in the genome of the intended host cell in proximity and orientation sufficient to influence the transcription of DNA encoding the desired IL-1ra-L polypeptide. The control element controls a portion of the DNA present in the host cell genome. Thus, the expression of the desired IL-1ra-L polypeptide may be achieved not by transfection of DNA that encodes the IL-1ra-L gene itself, but rather by the use of targeting DNA (containing regions of homology with the endogenous gene of interest) coupled with DNA regulatory segments that provide the endogenous gene sequence with recognizable signals for transcription of an IL-1ra-L gene.

In an exemplary method, the expression of a desired targeted gene in a cell (*i.e.*, a desired endogenous cellular gene) is altered via homologous recombination into the cellular genome at a preselected site, by the introduction of DNA which includes at least a regulatory sequence, an exon, and a splice donor site. These components are introduced into the chromosomal (genomic) DNA in such a manner that this, in effect, results in the production of a new transcription unit (in which the regulatory sequence, the exon, and the splice donor site present in the DNA construct are operatively linked to the endogenous gene). As a result of the introduction of these components into the chromosomal DNA, the expression of the desired endogenous gene is altered.

Altered gene expression, as described herein, encompasses activating (or causing to be expressed) a gene which is normally silent (unexpressed) in the cell as obtained, as well as increasing the expression of a gene which is not expressed at physiologically significant levels in the cell as obtained. The embodiments further encompass changing the pattern of regulation or induction such that it is different from the pattern of regulation or induction that occurs in the cell as obtained, and reducing (including eliminating) the expression of a gene which is expressed in the cell as obtained.

One method by which homologous recombination can be used to increase, or cause, IL-1ra-L polypeptide production from a cell's endogenous IL-1ra-L gene involves first using homologous recombination to place a recombination sequence from a site-specific recombination system (*e.g.*, Cre/loxP, FLP/FRT) (Sauer, 1994, *Curr. Opin. Biotechnol.*, 5:521-27; Sauer, 1993, *Methods Enzymol.*, 225:890-900) upstream of (*i.e.*, 5' to) the cell's endogenous genomic IL-1ra-L polypeptide coding region. A plasmid containing a recombination site homologous to the site that was placed just upstream of the genomic IL-1ra-L polypeptide coding region is introduced into the modified cell line along with the appropriate recombinase enzyme. This recombinase causes the plasmid to integrate, via the plasmid's recombination site, into the recombination site located just upstream of the genomic IL-1ra-L polypeptide coding region in the cell line (Baubonis and Sauer, 1993, *Nucleic Acids Res.* 21:2025-29; O'Gorman *et al.*, 1991, *Science* 251:1351-55). Any flanking sequences known to increase transcription (*e.g.*, enhancer/promoter, intron, translational enhancer), if properly positioned in this plasmid, would integrate in such a manner as to create a new or modified transcriptional unit resulting in *de novo* or increased IL-1ra-L polypeptide production from the cell's endogenous IL-1ra-L gene.

A further method to use the cell line in which the site specific recombination sequence had been placed just upstream of the cell's endogenous genomic IL-1ra-L polypeptide coding region is to use homologous recombination to introduce a second recombination site elsewhere in the cell line's genome. The appropriate recombinase enzyme is then introduced into the two-recombination-site cell line, causing a recombination event (deletion, inversion, and translocation) (Sauer, 1994, *Curr. Opin. Biotechnol.*, 5:521-27; Sauer, 1993, *Methods Enzymol.*, 225:890-900) that would create a new or modified transcriptional unit resulting in *de novo* or increased IL-1ra-L polypeptide production from the cell's endogenous IL-1ra-L gene.

An additional approach for increasing, or causing, the expression of IL-1ra-L polypeptide from a cell's endogenous IL-1ra-L gene involves increasing, or causing, the expression of a gene or genes (*e.g.*, transcription factors) and/or

decreasing the expression of a gene or genes (*e.g.*, transcriptional repressors) in a manner which results in *de novo* or increased IL-1ra-L polypeptide production from the cell's endogenous IL-1ra-L gene. This method includes the introduction of a non-naturally occurring polypeptide (*e.g.*, a polypeptide comprising a site  
5 specific DNA binding domain fused to a transcriptional factor domain) into the cell such that *de novo* or increased IL-1ra-L polypeptide production from the cell's endogenous IL-1ra-L gene results.

The present invention further relates to DNA constructs useful in the method of altering expression of a target gene. In certain embodiments, the  
10 exemplary DNA constructs comprise: (a) one or more targeting sequences, (b) a regulatory sequence, (c) an exon, and (d) an unpaired splice-donor site. The targeting sequence in the DNA construct directs the integration of elements (a) - (d) into a target gene in a cell such that the elements (b) - (d) are operatively linked to sequences of the endogenous target gene. In another embodiment, the  
15 DNA constructs comprise: (a) one or more targeting sequences, (b) a regulatory sequence, (c) an exon, (d) a splice-donor site, (e) an intron, and (f) a splice-acceptor site, wherein the targeting sequence directs the integration of elements (a) - (f) such that the elements of (b) - (f) are operatively linked to the endogenous gene. The targeting sequence is homologous to the preselected site in the cellular  
20 chromosomal DNA with which homologous recombination is to occur. In the construct, the exon is generally 3' of the regulatory sequence and the splice-donor site is 3' of the exon.

If the sequence of a particular gene is known, such as the nucleic acid sequence of IL-1ra-L polypeptide presented herein, a piece of DNA that is  
25 complementary to a selected region of the gene can be synthesized or otherwise obtained, such as by appropriate restriction of the native DNA at specific recognition sites bounding the region of interest. This piece serves as a targeting sequence upon insertion into the cell and will hybridize to its homologous region within the genome. If this hybridization occurs during DNA replication, this piece  
30 of DNA, and any additional sequence attached thereto, will act as an Okazaki fragment and will be incorporated into the newly synthesized daughter strand of



DNA. The present invention, therefore, includes nucleotides encoding an IL-1ra-L polypeptide, which nucleotides may be used as targeting sequences.

IL-1ra-L polypeptide cell therapy, *e.g.*, the implantation of cells producing IL-1ra-L polypeptides, is also contemplated. This embodiment involves  
5 implanting cells capable of synthesizing and secreting a biologically active form of IL-1ra-L polypeptide. Such IL-1ra-L polypeptide-producing cells can be cells that are natural producers of IL-1ra-L polypeptides or may be recombinant cells whose ability to produce IL-1ra-L polypeptides has been augmented by transformation with a gene encoding the desired IL-1ra-L polypeptide or with a  
10 gene augmenting the expression of IL-1ra-L polypeptide. Such a modification may be accomplished by means of a vector suitable for delivering the gene as well as promoting its expression and secretion. In order to minimize a potential immunological reaction in patients being administered an IL-1ra-L polypeptide, as may occur with the administration of a polypeptide of a foreign species, it is  
15 preferred that the natural cells producing IL-1ra-L polypeptide be of human origin and produce human IL-1ra-L polypeptide. Likewise, it is preferred that the recombinant cells producing IL-1ra-L polypeptide be transformed with an expression vector containing a gene encoding a human IL-1ra-L polypeptide.

Implanted cells may be encapsulated to avoid the infiltration of  
20 surrounding tissue. Human or non-human animal cells may be implanted in patients in biocompatible, semipermeable polymeric enclosures or membranes that allow the release of IL-1ra-L polypeptide, but that prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissue. Alternatively, the patient's own cells, transformed to produce  
25 IL-1ra-L polypeptides *ex vivo*, may be implanted directly into the patient without such encapsulation.

Techniques for the encapsulation of living cells are known in the art, and the preparation of the encapsulated cells and their implantation in patients may be routinely accomplished. For example, Baetge *et al.* (PCT Pub. No. WO 95/05452  
30 and PCT/US94/09299) describe membrane capsules containing genetically engineered cells for the effective delivery of biologically active molecules. The

capsules are biocompatible and are easily retrievable. The capsules encapsulate cells transfected with recombinant DNA molecules comprising DNA sequences coding for biologically active molecules operatively linked to promoters that are not subject to down-regulation *in vivo* upon implantation into a mammalian host.

5 The devices provide for the delivery of the molecules from living cells to specific sites within a recipient. In addition, *see* U.S. Patent Nos. 4,892,538; 5,011,472; and 5,106,627. A system for encapsulating living cells is described in PCT Pub. No. WO 91/10425 (Aebischer *et al.*). *See also*, PCT Pub. No. WO 91/10470 (Aebischer *et al.*); Winn *et al.*, 1991, *Exper. Neurol.* 113:322-29; Aebischer *et al.*,  
10 1991, *Exper. Neurol.* 111:269-75; and Tresco *et al.*, 1992, *ASAIO* 38:17-23.

*In vivo* and *in vitro* gene therapy delivery of IL-1ra-L polypeptides is also envisioned. One example of a gene therapy technique is to use the IL-1ra-L gene (either genomic DNA, cDNA, and/or synthetic DNA) encoding an IL-1ra-L polypeptide which may be operably linked to a constitutive or inducible promoter  
15 to form a "gene therapy DNA construct." The promoter may be homologous or heterologous to the endogenous IL-1ra-L gene, provided that it is active in the cell or tissue type into which the construct will be inserted. Other components of the gene therapy DNA construct may optionally include DNA molecules designed for site-specific integration (*e.g.*, endogenous sequences useful for homologous  
20 recombination), tissue-specific promoters, enhancers or silencers, DNA molecules capable of providing a selective advantage over the parent cell, DNA molecules useful as labels to identify transformed cells, negative selection systems, cell specific binding agents (as, for example, for cell targeting), cell-specific internalization factors, transcription factors enhancing expression from a vector,  
25 and factors enabling vector production.

A gene therapy DNA construct can then be introduced into cells (either *ex vivo* or *in vivo*) using viral or non-viral vectors. One means for introducing the gene therapy DNA construct is by means of viral vectors as described herein. Certain vectors, such as retroviral vectors, will deliver the DNA construct to the  
30 chromosomal DNA of the cells, and the gene can integrate into the chromosomal DNA. Other vectors will function as episomes, and the gene therapy DNA

construct will remain in the cytoplasm.

In yet other embodiments, regulatory elements can be included for the controlled expression of the IL-1ra-L gene in the target cell. Such elements are turned on in response to an appropriate effector. In this way, a therapeutic  
5 polypeptide can be expressed when desired. One conventional control means involves the use of small molecule dimerizers or rapalogs to dimerize chimeric proteins which contain a small molecule-binding domain and a domain capable of initiating a biological process, such as a DNA-binding protein or transcriptional activation protein (*see* PCT Pub. Nos. WO 96/41865, WO 97/31898, and WO  
10 97/31899). The dimerization of the proteins can be used to initiate transcription of the transgene.

An alternative regulation technology uses a method of storing proteins expressed from the gene of interest inside the cell as an aggregate or cluster. The gene of interest is expressed as a fusion protein that includes a conditional  
15 aggregation domain that results in the retention of the aggregated protein in the endoplasmic reticulum. The stored proteins are stable and inactive inside the cell. The proteins can be released, however, by administering a drug (*e.g.*, small molecule ligand) that removes the conditional aggregation domain and thereby specifically breaks apart the aggregates or clusters so that the proteins may be  
20 secreted from the cell. *See Aridor et al.*, 2000, *Science* 287:816-17 and *Rivera et al.*, 2000, *Science* 287:826-30.

Other suitable control means or gene switches include, but are not limited to, the systems described herein. Mifepristone (RU486) is used as a progesterone antagonist. The binding of a modified progesterone receptor ligand-binding  
25 domain to the progesterone antagonist activates transcription by forming a dimer of two transcription factors that then pass into the nucleus to bind DNA. The ligand-binding domain is modified to eliminate the ability of the receptor to bind to the natural ligand. The modified steroid hormone receptor system is further described in U.S. Patent No. 5,364,791 and PCT Pub. Nos. WO 96/40911 and  
30 WO 97/10337.

Yet another control system uses ecdysone (a fruit fly steroid hormone) which binds to and activates an ecdysone receptor (cytoplasmic receptor). The receptor then translocates to the nucleus to bind a specific DNA response element (promoter from ecdysone-responsive gene). The ecdysone receptor includes a  
5 transactivation domain, DNA-binding domain, and ligand-binding domain to initiate transcription. The ecdysone system is further described in U.S. Patent No. 5,514,578 and PCT Pub. Nos. WO 97/38117, WO 96/37609, and WO 93/03162.

Another control means uses a positive tetracycline-controllable transactivator. This system involves a mutated tet repressor protein DNA-binding  
10 domain (mutated tet R-4 amino acid changes which resulted in a reverse tetracycline-regulated transactivator protein, *i.e.*, it binds to a tet operator in the presence of tetracycline) linked to a polypeptide which activates transcription. Such systems are described in U.S. Patent Nos. 5,464,758, 5,650,298, and 5,654,168.

15 Additional expression control systems and nucleic acid constructs are described in U.S. Patent Nos. 5,741,679 and 5,834,186, to Innovir Laboratories Inc.

*In vivo* gene therapy may be accomplished by introducing the gene encoding IL-1ra-L polypeptide into cells via local injection of an IL-1ra-L nucleic  
20 acid molecule or by other appropriate viral or non-viral delivery vectors. Hefti, 1994, *Neurobiology* 25:1418-35. For example, a nucleic acid molecule encoding an IL-1ra-L polypeptide may be contained in an adeno-associated virus (AAV) vector for delivery to the targeted cells (*see, e.g.*, Johnson, PCT Pub. No. WO 95/34670; PCT App. No. PCT/US95/07178). The recombinant AAV genome  
25 typically contains AAV inverted terminal repeats flanking a DNA sequence encoding an IL-1ra-L polypeptide operably linked to functional promoter and polyadenylation sequences.

Alternative suitable viral vectors include, but are not limited to, retrovirus, adenovirus, herpes simplex virus, lentivirus, hepatitis virus, parvovirus,  
30 papovavirus, poxvirus, alphavirus, coronavirus, rhabdovirus, paramyxovirus, and papilloma virus vectors. U.S. Patent No. 5,672,344 describes an *in vivo* viral-

mediated gene transfer system involving a recombinant neurotrophic HSV-1 vector. U.S. Patent No. 5,399,346 provides examples of a process for providing a patient with a therapeutic protein by the delivery of human cells which have been treated *in vitro* to insert a DNA segment encoding a therapeutic protein.

5 Additional methods and materials for the practice of gene therapy techniques are described in U.S. Patent Nos. 5,631,236 (involving adenoviral vectors), 5,672,510 (involving retroviral vectors), 5,635,399 (involving retroviral vectors expressing cytokines).

Nonviral delivery methods include, but are not limited to, liposome-mediated transfer, naked DNA delivery (direct injection), receptor-mediated transfer (ligand-DNA complex), electroporation, calcium phosphate precipitation, and microparticle bombardment (*e.g.*, gene gun). Gene therapy materials and methods may also include inducible promoters, tissue-specific enhancer-promoters, DNA sequences designed for site-specific integration, DNA sequences  
15 capable of providing a selective advantage over the parent cell, labels to identify transformed cells, negative selection systems and expression control systems (safety measures), cell-specific binding agents (for cell targeting), cell-specific internalization factors, and transcription factors to enhance expression by a vector as well as methods of vector manufacture. Such additional methods and materials  
20 for the practice of gene therapy techniques are described in U.S. Patent Nos. 4,970,154 (involving electroporation techniques), 5,679,559 (describing a lipoprotein-containing system for gene delivery), 5,676,954 (involving liposome carriers), 5,593,875 (describing methods for calcium phosphate transfection), and 4,945,050 (describing a process wherein biologically active particles are propelled  
25 at cells at a speed whereby the particles penetrate the surface of the cells and become incorporated into the interior of the cells), and PCT Pub. No. WO 96/40958 (involving nuclear ligands).

It is also contemplated that IL-1ra-L gene therapy or cell therapy can further include the delivery of one or more additional polypeptide(s) in the same  
30 or a different cell(s). Such cells may be separately introduced into the patient, or the cells may be contained in a single implantable device, such as the

encapsulating membrane described above, or the cells may be separately modified by means of viral vectors.

A means to increase endogenous IL-1ra-L polypeptide expression in a cell via gene therapy is to insert one or more enhancer elements into the IL-1ra-L polypeptide promoter, where the enhancer elements can serve to increase transcriptional activity of the IL-1ra-L gene. The enhancer elements used will be selected based on the tissue in which one desires to activate the gene – enhancer elements known to confer promoter activation in that tissue will be selected. For example, if a gene encoding an IL-1ra-L polypeptide is to be “turned on” in T-cells, the *lck* promoter enhancer element may be used. Here, the functional portion of the transcriptional element to be added may be inserted into a fragment of DNA containing the IL-1ra-L polypeptide promoter (and optionally, inserted into a vector and/or 5’ and/or 3’ flanking sequences) using standard cloning techniques. This construct, known as a “homologous recombination construct,” can then be introduced into the desired cells either *ex vivo* or *in vivo*.

Gene therapy also can be used to decrease IL-1ra-L polypeptide expression by modifying the nucleotide sequence of the endogenous promoter. Such modification is typically accomplished via homologous recombination methods. For example, a DNA molecule containing all or a portion of the promoter of the IL-1ra-L gene selected for inactivation can be engineered to remove and/or replace pieces of the promoter that regulate transcription. For example, the TATA box and/or the binding site of a transcriptional activator of the promoter may be deleted using standard molecular biology techniques; such deletion can inhibit promoter activity thereby repressing the transcription of the corresponding IL-1ra-L gene. The deletion of the TATA box or the transcription activator binding site in the promoter may be accomplished by generating a DNA construct comprising all or the relevant portion of the IL-1ra-L polypeptide promoter (from the same or a related species as the IL-1ra-L gene to be regulated) in which one or more of the TATA box and/or transcriptional activator binding site nucleotides are mutated via substitution, deletion and/or insertion of one or more nucleotides. As a result, the TATA box and/or activator binding site has

decreased activity or is rendered completely inactive. This construct, which also will typically contain at least about 500 bases of DNA that correspond to the native (endogenous) 5' and 3' DNA sequences adjacent to the promoter segment that has been modified, may be introduced into the appropriate cells (either *ex vivo* or *in vivo*) either directly or via a viral vector as described herein. Typically, the integration of the construct into the genomic DNA of the cells will be via homologous recombination, where the 5' and 3' DNA sequences in the promoter construct can serve to help integrate the modified promoter region via hybridization to the endogenous chromosomal DNA.

10

#### Therapeutic Uses

IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists thereof can be used to treat, diagnose, ameliorate, or prevent a number of diseases, disorders, or conditions, including those recited herein.

15

IL-1ra-L polypeptide agonists and antagonists include those molecules which regulate IL-1ra-L polypeptide activity and either increase or decrease at least one activity of the mature form of the IL-1ra-L polypeptide. Agonists or antagonists may be co-factors, such as a protein, peptide, carbohydrate, lipid, or small molecular weight molecule, which interact with IL-1ra-L polypeptide and thereby regulate its activity. Potential polypeptide agonists or antagonists include antibodies that react with either soluble or membrane-bound forms of IL-1ra-L polypeptides that comprise part or all of the extracellular domains of the said proteins. Molecules that regulate IL-1ra-L polypeptide expression typically include nucleic acids encoding IL-1ra-L polypeptide that can act as anti-sense regulators of expression.

20  
25

For example, the IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can be used to treat, diagnose, ameliorate, or prevent diseases, disorders, or conditions involving immune system dysfunction. Examples of such diseases include, but are not limited to, rheumatoid arthritis, psoriatic arthritis, inflammatory arthritis, osteoarthritis, inflammatory joint disease, autoimmune disease (including autoimmune

30

vasculitis), multiple sclerosis, lupus, diabetes (*e.g.*, insulin diabetes), inflammatory bowel disease, transplant rejection, graft versus host disease, and inflammatory conditions resulting from strain, sprain, cartilage damage, trauma, orthopedic surgery, infection or other disease processes. Other diseases  
5 influenced by the dysfunction of the immune system are encompassed within the scope of the invention.

The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat, diagnose, ameliorate, or prevent diseases, disorders, or conditions involving infection. Examples of such  
10 diseases include, but are not limited to, leprosy, viral infections (such as hepatitis or HIV), bacterial infection (such as *clostridium*-associated illnesses, including *clostridium*-associated diarrhea), pulmonary tuberculosis, acute febrile illness, fever, acute phase response of the liver, septicemia, or septic shock. Other diseases involving infection are encompassed within the scope of the invention.

15 The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat, diagnose, ameliorate, or prevent diseases, disorders, or conditions involving weight disorders. Examples of such diseases include, but are not limited to obesity, anorexia, cachexia (including AIDS-induced cachexia), myopathies (*e.g.*, muscle protein metabolism,  
20 such as in sepsis), and hypoglycemia. Other diseases involving weight disorders are encompassed within the scope of the invention.

The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat, diagnose, ameliorate, or prevent diseases, disorders, or conditions involving neuronal dysfunction.  
25 Examples of such diseases include, but are not limited to, Alzheimer's disease, Parkinson's disease, neurotoxicity (*e.g.*, as induced by HIV), ALS, brain injury, stress, depression, nociception and other pain (including cancer-related pain), hyperalgesia, epilepsy, learning impairment and memory disorders, sleep disturbance, and peripheral and central neuropathies. Other neurological  
30 disorders are encompassed within the scope of the invention.



The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat, diagnose, ameliorate, or prevent diseases, disorders, or conditions involving the lung. Examples of such diseases include, but are not limited to, acute or chronic lung injury (including  
5 interstitial lung disease), acute respiratory disease syndrome, pulmonary hypertension, emphysema, cystic fibrosis, pulmonary fibrosis, and asthma. Other diseases of the lung are encompassed within the scope of the invention.

The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat, diagnose, ameliorate, or  
10 prevent diseases, disorders, or conditions involving the skin. Examples of such diseases include, but are not limited to, psoriasis, eczema, and wound healing. Other diseases of the skin are encompassed within the scope of the invention.

The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat, diagnose, ameliorate, or  
15 prevent diseases, disorders, or conditions involving the kidney. Examples of such diseases include, but are not limited to, acute and chronic glomerulonephritis. Other diseases of the kidney are encompassed within the scope of the invention.

The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat, diagnose, ameliorate, or  
20 prevent diseases, disorders, or conditions involving the bone. Examples of such diseases include, but are not limited to, osteoporosis, osteopetrosis, osteogenesis imperfecta, Paget's disease, periodontal disease, temporal mandibular joint disease, and hypercalcemia. Other diseases of the bone are encompassed within the scope of the invention.

25 The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat, diagnose, ameliorate, or prevent diseases, disorders, or conditions involving the vascular system. Examples of such diseases include, but are not limited to, hemorrhage or stroke, hemorrhagic shock, ischemia (including cardiac ischemia and cerebral ischemia,  
30 *e.g.*, brain injury as a result of trauma, epilepsy, hemorrhage or stroke, each of which may lead to neurodegeneration), atherosclerosis, congestive heart failure,

restenosis, reperfusion injury, and angiogenesis. Other diseases of the vascular system are encompassed within the scope of the invention.

The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat, diagnose, ameliorate, or prevent diseases, disorders, or conditions involving tumor cells. Examples of such diseases include, but are not limited to, lymphomas, bone sarcoma, chronic and acute myelogenous leukemia (CML and AML) and other leukemias, multiple myeloma, lung cancer, breast cancer, tumor metastasis, and side effects from radiation therapy. Other diseases involving tumor cells are encompassed within the scope of the invention.

The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat, diagnose, ameliorate, or prevent diseases, disorders, or conditions involving the reproductive system. Examples of such diseases include, but are not limited to, infertility, miscarriage, pre-term labor and delivery, and endometriosis. Other diseases involving the reproductive system are encompassed within the scope of the invention.

The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat, diagnose, ameliorate, or prevent diseases, disorders, or conditions involving the eye. Examples of such diseases include, but are not limited to, inflammatory eye disease (as may be associated with, for example, corneal transplant), retinal degeneration, blindness, macular degeneration, glaucoma, uveitis, and retinal neuropathy. Other diseases of the eye are encompassed within the scope of the invention.

The IL-1ra-L nucleic acid molecules, polypeptides, and agonists and antagonists of the invention can also be used to treat diseases such as acute pancreatitis, chronic fatigue syndrome, fibromyalgia, and Kawasaki's disease (MLNS).

IL-1 inhibitors include any protein capable of specifically preventing activation of cellular receptors to IL-1, which may result from any number of mechanisms. Such mechanisms include down-regulating IL-1 production, binding free IL-1, interfering with IL-1 binding to its receptor, interfering with the

formation of the IL-1 receptor complex (*i.e.*, association of IL-1 receptor with IL-1 receptor accessory protein), and interfering with the modulation of IL-1 signaling after binding to its receptor. Such interleukin-1 inhibitors include: interleukin-1 receptor antagonists such as IL-1ra-L, as described herein, anti-IL-1  
5 receptor monoclonal antibodies (*e.g.*, European Patent No. 623674), IL-1 binding proteins such as soluble IL-1 receptors (*e.g.*, U.S. Patent Nos. 5,492,888, 5,488,032, 5,464,937, 5,319,071, and 5,180,812), anti-IL-1 monoclonal antibodies (*e.g.*, PCT Pub. Nos. WO 95/01997, WO 94/02627, WO 90/06371; U.S. Patent No. 4,935,343; and European Patent Nos. 364778, 267611 and 220063), IL-1  
10 receptor accessory proteins and antibodies thereto (*e.g.*, PCT Pub. No. WO 96/23067); inhibitors of interleukin-1 $\beta$  converting enzyme (ICE) or caspase I, which can be used to inhibit IL-1 $\beta$  production and secretion, interleukin-1 $\beta$  protease inhibitors, and other compounds and proteins which block *in vivo* synthesis or extracellular release of IL-1.

15 Exemplary IL-1 inhibitors are disclosed in US Patent Nos. 5,747,444, 5,359,032, 5,608,035, 5,843,905, 5,359,032, 5,866,576, 5,869,660, 5,869,315, 5,872,095, 5,955,480; PCT Pub. Nos. WO 98/21957, WO 96/09323, WO 91/17184, WO 96/40907, WO 98/32733, WO 98/42325, WO 98/44940, WO 98/47892, WO 98/56377, WO 99/03837, WO 99/06426, WO 99/06042, WO  
20 91/17249, WO 98/32733, WO 98/17661, WO 97/08174, WO 95/34326, WO 99/36426, and WO 99/36415; European Patent Nos. 534978 and 894795; and French Patent Application FR 2762514.

Interleukin-1 receptor antagonist (IL-1ra) is a human protein that acts as a natural inhibitor of interleukin-1. Preferred receptor antagonists (including IL-1ra  
25 and variants and derivatives thereof), as well as methods of making and using thereof, are described in U.S. Patent No. 5,075,222; PCT Pub. Nos. WO 91/08285, WO 91/17184, WO 92/16221, WO 93/21946, WO 94/06457, WO 94/21275, WO 94/21235, WO 94/20517, WO 96/22793, WO 97/28828, and WO 99/36541; Austrian Patent No. AU 9173636; French Patent No. FR 2706772; and  
30 German Patent No. DE 4219626. Such proteins include glycosylated as well as non-glycosylated IL-1 receptor antagonists.

Specifically, three exemplary forms of IL-1ra and variants thereof are disclosed and described in the 5,075,222 patent. The first of these, called "IL-1i," is characterized as a 22-23 kD molecule on SDS-PAGE with an approximate isoelectric point of 4.8, eluting from a MonoQ FPLC column at around 52 mM NaCl in Tris buffer, pH 7.6. The second, IL-1ra $\beta$ , is characterized as a 22-23 kD protein, eluting from a MonoQ column at 48 mM NaCl. Both IL-1ra $\alpha$  and IL-1ra $\beta$  are glycosylated. The third, IL-1ra $\chi$ , is characterized as a 20 kD protein, eluting from a MonoQ column at 48 mM NaCl, and is non-glycosylated. U.S. Patent No. 5,075,222 also discloses methods for isolating the genes responsible for coding the inhibitors, cloning the gene in suitable vectors and cell types, and expressing the gene to produce the inhibitors.

Those skilled in the art will recognize that many combinations of deletions, insertions, and substitutions (individually or collectively "variant(s)" herein) can be made within the amino acid sequences of IL-1ra-L, provided that the resulting molecule is biologically active (*e.g.*, possesses the ability to affect one or more of the diseases and disorders such as those recited herein.)

As contemplated by the present invention, an IL-1ra-L polypeptide may be administered as an adjunct to other therapy and also with other pharmaceutical compositions suitable for the indication being treated. An IL-1ra-L polypeptide and any of one or more additional therapies or pharmaceutical formulations may be administered separately, sequentially, or simultaneously.

In a specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pre-treatment, post-treatment, or concurrent treatment) with any of one or more TNF inhibitors for the treatment or prevention of the diseases and disorders recited herein.

Such TNF inhibitors include compounds and proteins that block *in vivo* synthesis or extracellular release of TNF. In a specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pre-treatment, post-treatment, or concurrent treatment) with any of one or more of the following TNF inhibitors: TNF binding proteins (soluble TNF receptor type-I and soluble TNF receptor type-II ("sTNFRs"), as defined herein), anti-TNF

antibodies, granulocyte colony stimulating factor, thalidomide, BN 50730, tenidap, E 5531, tiapafant PCA 4248, nimesulide, panavir, rolipram, RP 73401, peptide T, MDL 201,449A, (1R,3S)-Cis-1-[9-(2,6-diaminopuriny)]-3-hydroxy-4-cyclopentene hydrochloride, (1R,3R)-trans-1-(9-(2,6-diamino)purine]-3-acetoxycyclopentane, (1R,3R)-trans-1-[9-adenyl)-3-azidocyclopentane hydrochloride and (1R,3R)-trans-1-(6-hydroxy-purin-9-yl)-3-azidocyclopentane. TNF binding proteins are disclosed in the art (U.S. Patent No. 5,136,021; European Patent Nos. 308378, 422339, 393438, 398327, 412486, 418014, 433900, 464533, 512528, 526905, 568928, 417563; PCT Pub. Nos. WO 90/13575, WO 91/03553, WO 92/01002, WO 92/13095, WO 92/16221, WO 93/07863, WO 93/21946, WO 93/19777, WO 94/06476, PCT App. No. PCT/US97/12244; English Patent Nos. GB 2218101 and 2246569; and Japanese Patent App. No. JP 127,800/1991).

For example, European Patent Nos. 393438 and 422339 teach the amino acid and nucleic acid sequences of a soluble TNF receptor type I (also known as "sTNFR-I" or "30kDa TNF inhibitor") and a soluble TNF receptor type II (also known as "sTNFR-II" or "40kDa TNF inhibitor"), collectively termed "sTNFRs," as well as modified forms thereof (*e.g.*, fragments, functional derivatives, and variants). European Patent Nos. 393438 and 422339 also disclose methods for isolating the genes responsible for coding the inhibitors, cloning the gene in suitable vectors and cell types, and expressing the gene to produce the inhibitors. Additionally, polyvalent forms (*i.e.*, molecules comprising more than one active moiety) of sTNFR-I and sTNFR-II have also been disclosed. In one embodiment, the polyvalent form may be constructed by chemically coupling at least one TNF inhibitor and another moiety with any clinically acceptable linker, for example polyethylene glycol (PCT Pub. Nos. WO 92/16221 and WO 95/34326), by a peptide linker (Neve *et al.*, 1996, *Cytokine*, 8:365-70), by chemically coupling to biotin and then binding to avidin (PCT Pub. No. WO 91/03553) and, finally, by combining chimeric antibody molecules (U.S. Patent No. 5,116,964; PCT Pub. Nos. WO 89/09622 and WO 91/16437; and European Patent No. 315062).

Anti-TNF antibodies include MAK 195F Fab antibody (Holler *et al.*,

1993, *1st International Symposium on Cytokines in Bone Marrow Transplantation* 147), CDP 571 anti-TNF monoclonal antibody (Rankin *et al.*, 1995, *Br. J. Rheumatol.*, 34:334-42), BAY X 1351 murine anti-tumor necrosis factor monoclonal antibody (Kieft *et al.*, 1995, *7th European Congress of Clinical*  
5 *Microbiology and Infectious Diseases* 9); CenTNF cA2 anti-TNF monoclonal antibody (Elliott *et al.*, 1994, *Lancet*, 344:1125-27; Elliott *et al.*, 1994, *Lancet*, 344:1105-10).

In a specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment, or concurrent  
10 treatment) with secreted or soluble human fas antigen or recombinant versions thereof (PCT Pub. No. WO 96/20206; Mountz *et al.*, 1995, *J. Immunol.*, 155:4829-37; and European Patent No. 510691). PCT Pub. No. WO 96/20206 discloses secreted human fas antigen (native and recombinant, including an Ig fusion protein), methods for isolating the genes responsible for coding the soluble  
15 recombinant human fas antigen, methods for cloning the gene in suitable vectors and cell types, and methods for expressing the gene to produce the inhibitors. European Patent No. 510691 teaches nucleic acids coding for human fas antigen, including soluble fas antigen, vectors expressing for said nucleic acids, and transformants transfected with the vector. When administered parenterally, doses  
20 of a secreted or soluble fas antigen fusion protein each are generally from about 1 µg/kg to about 100 µg/kg.

Current treatment of the diseases and disorders recited herein, including acute and chronic inflammation such as rheumatic diseases, commonly includes the use of first line drugs for control of pain and inflammation; these drugs are  
25 classified as non-steroidal, anti-inflammatory drugs (NSAIDs). Secondary treatments include corticosteroids, slow acting antirheumatic drugs (SAARDs), or disease modifying (DM) drugs. Information regarding the following compounds can be found in *The Merck Manual of Diagnosis and Therapy* (16th ed. 1992) and in *Pharmaprojects* (PJB Publications Ltd).

30 In a specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide and any of one or more NSAIDs for the treatment of the

diseases and disorders recited herein, including acute and chronic inflammation such as rheumatic diseases, and graft versus host disease. NSAIDs owe their anti-inflammatory action, at least in part, to the inhibition of prostaglandin synthesis (Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (7th ed.

5    1985)). NSAIDs can be characterized into at least nine groups: (1) salicylic acid derivatives, (2) propionic acid derivatives, (3) acetic acid derivatives, (4) fenamic acid derivatives, (5) carboxylic acid derivatives, (6) butyric acid derivatives, (7) oxicams, (8) pyrazoles, and (9) pyrazolones.

10    In another specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment, or concurrent treatment) with any of one or more salicylic acid derivatives, prodrug esters, or pharmaceutically acceptable salts thereof. Such salicylic acid derivatives, prodrug esters, and pharmaceutically acceptable salts thereof comprise: acetaminosalol, aloxiprin, aspirin, benorylate, bromosaligenin, calcium  
15    acetylsalicylate, choline magnesium trisalicylate, magnesium salicylate, choline salicylate, diflusal, etersalate, fendosal, gentisic acid, glycol salicylate, imidazole salicylate, lysine acetylsalicylate, mesalamine, morpholine salicylate, 1-naphthyl salicylate, olsalazine, parsalimide, phenyl acetylsalicylate, phenyl salicylate, salacetamide, salicylamide O-acetic acid, salsalate, sodium salicylate  
20    and sulfasalazine. Structurally related salicylic acid derivatives having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

25    In an additional specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment, or concurrent treatment) with any of one or more propionic acid derivatives, prodrug esters, or pharmaceutically acceptable salts thereof. The propionic acid derivatives, prodrug esters, and pharmaceutically acceptable salts thereof comprise: alminoprofen, benoxaprofen, bucloxic acid, carprofen, dexindoprofen, fenoprofen, flunoxaprofen, fluprofen, flurbiprofen, furocloprofen, ibuprofen,  
30    ibuprofen aluminum, ibuprofen, indoprofen, isoprofen, ketoprofen, loxoprofen, mioprofen, naproxen, naproxen sodium, oxaprozin, piketoprofen, pimeprofen,

pirprofen, pranoprofen, protizinic acid, pyridoxiprofen, suprofen, tiaprofenic acid and tioxaprofen. Structurally related propionic acid derivatives having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

5 In yet another specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment, or concurrent treatment) with any of one or more acetic acid derivatives, prodrug esters, or pharmaceutically acceptable salts thereof. The acetic acid derivatives, prodrug esters, and pharmaceutically acceptable salts thereof comprise:  
10 acemetacin, alclofenac, amfenac, bufexamac, cinmetacin, clopirac, delmetacin, diclofenac potassium, diclofenac sodium, etodolac, felbinac, fenclofenac, fenclorac, fenclozic acid, fentiazac, furofenac, glucametacin, ibufenac, indomethacin, isofezolac, isoxepac, lonazolac, metiazinic acid, oxametacin, oxpinac, pimetacin, proglumetacin, sulindac, talmetacin, tiaramide, tiopinac,  
15 tolmetin, tolmetin sodium, zidometacin and zomepirac. Structurally related acetic acid derivatives having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

In another specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment, or  
20 concurrent treatment) with any of one or more fenamic acid derivatives, prodrug esters, or pharmaceutically acceptable salts thereof. The fenamic acid derivatives, prodrug esters, and pharmaceutically acceptable salts thereof comprise: enfenamic acid, etofenamate, flufenamic acid, isonixin, meclofenamic acid, meclofenamate sodium, medofenamic acid, mefenamic acid, niflumic acid, talniflumate,  
25 terofenamate, tolfenamic acid and ufenamate. Structurally related fenamic acid derivatives having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

In an additional specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment,  
30 or concurrent treatment) with any of one or more carboxylic acid derivatives, prodrug esters, or pharmaceutically acceptable salts thereof. The carboxylic acid



derivatives, prodrug esters, and pharmaceutically acceptable salts thereof which can be used comprise: clidanac, diflunisal, flufenisal, inoridine, ketorolac and tinoridine. Structurally related carboxylic acid derivatives having similar analgesic and anti-inflammatory properties are also intended to be encompassed  
5 by this group.

In yet another specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment, or concurrent treatment) with any of one or more butyric acid derivatives, prodrug esters, or pharmaceutically acceptable salts thereof. The butyric acid derivatives,  
10 prodrug esters, and pharmaceutically acceptable salts thereof comprise: bumadizon, butibufen, fenbufen and xenbucin. Structurally related butyric acid derivatives having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

In another specific embodiment, the present invention is directed to the use  
15 of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment, or concurrent treatment) with any of one or more oxicams, prodrug esters, or pharmaceutically acceptable salts thereof. The oxicams, prodrug esters, and pharmaceutically acceptable salts thereof comprise: droxicam, enolicam, isoxicam, piroxicam, sudoxicam, tenoxicam and 4-hydroxyl-1,2-benzothiazine  
20 1,1-dioxide 4-(N-phenyl)-carboxamide. Structurally related oxicams having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

In still another specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment,  
25 or concurrent treatment) with any of one or more pyrazoles, prodrug esters, or pharmaceutically acceptable salts thereof. The pyrazoles, prodrug esters, and pharmaceutically acceptable salts thereof which may be used comprise: difenamizole and epirizole. Structurally related pyrazoles having similar analgesic and anti-inflammatory properties are also intended to be encompassed  
30 by this group.

In an additional specific embodiment, the present invention is directed to

the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment or, concurrent treatment) with any of one or more pyrazolones, prodrug esters, or pharmaceutically acceptable salts thereof. The pyrazolones, prodrug esters, and pharmaceutically acceptable salts thereof which may be used comprise: apazone, azapropazone, benzpiperylon, feprazone, mofebutazone, morazone, oxyphenbutazone, phenylbutazone, pipebuzone, propylphenazone, ramifenazone, suxibuzone and thiazolinobutazone. Structurally related pyrazalones having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

10 In another specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment, or concurrent treatment) with any of one or more of the following: NSAIDs:  $\epsilon$ -acetamidocaproic acid, S-adenosylmethionine, 3-amino-4-hydroxybutyric acid, amixetrine, anitrazafen, antrafenine, bendazac, bendazac lysinate, benzydamine, beprozin, broperamole, bucolome, bufezolac, ciproquazone, cloximate, dazidamine, deboxamet, detomidine, difenpiramide, difenpyramide, difisalamine, ditazol, emorfazone, fanetizole mesylate, fenflumizole, floctafenine, flumizole, flunixin, fluproquazone, fopirtoline, fosfosal, guaimesal, guaiazolene, isonixirn, lefetamine HCl, leflunomide, lofemizole, lotifazole, lysin clonixinate, meseclazone, nabumetone, nictindole, nimesulide, orgotein, orpanoxin, oxaceprol, oxapadol, paranyline, perisoxal, perisoxal citrate, pifoxime, piroxten, pirazolac, 15 pirfenidone, proquazone, proxazole, thielavin B, tiiflamizole, timegadine, tolectin, tolpadol, tryptamid and those designated by company code number such as 480156S, AA861, AD1590, AFP802, AFP860, AI77B, AP504, AU8001, BPPC, BW540C, CHINOIN 127, CN100, EB382, EL508, F1044, FK-506, GV3658, ITF182, KCNTEI6090, KME4, LA2851, MR714, MR897, MY309, ONO3144, PR823, PV102, PV108, R830, RS2131, SCR152, SH440, SIR133, SPAS510, SQ27239, ST281, SY6001, TA60, TAI-901 (4-benzoyl-1-indancarboxylic acid), TVX2706, U60257, UR2301 and WY41770. Structurally related NSAIDs having 20 similar analgesic and anti-inflammatory properties to the NSAIDs are also intended to be encompassed by this group.

30

In still another specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment or concurrent treatment) with any of one or more corticosteroids, prodrug esters, or pharmaceutically acceptable salts thereof for the treatment of the diseases and disorders recited herein, including acute and chronic inflammation such as rheumatic diseases, graft versus host disease, and multiple sclerosis. Corticosteroids, prodrug esters, and pharmaceutically acceptable salts thereof include hydrocortisone and compounds which are derived from hydrocortisone, such as 21-acetoxypregnenolone, alclomerasone, algestone, amcinonide, beclomethasone, betamethasone, betamethasone valerate, budesonide, chloroprednisone, clobetasol, clobetasol propionate, clobetasone, clobetasone butyrate, clocortolone, cloprednol, corticosterone, cortisone, cortivazol, deflazacort, desonide, desoximetasone, dexamethasone, diflorasone, diflucortolone, difluprednate, enoxolone, fluazacort, flucoronide, flumethasone, flumethasone pivalate, flucinolone acetonide, flunisolide, fluocinonide, fluorocinolone acetonide, fluocortin butyl, fluocortolone, fluocortolone hexanoate, diflucortolone valerate, fluorometholone, fluperolone acetate, fluprednidene acetate, fluprednisolone, flurandrenolide, formocortol, halcinonide, halometasone, halopredone acetate, hydrocortamate, hydrocortisone, hydrocortisone acetate, hydrocortisone butyrate, hydrocortisone phosphate, hydrocortisone 21-sodium succinate, hydrocortisone tebutate, mазipredone, medrysone, meprednisone, methylprednisolone, mometasone furoate, paramethasone, prednicarbate, prednisolone, prednisolone 21-diedryaminoacetate, prednisolone sodium phosphate, prednisolone sodium succinate, prednisolone sodium 21-*m*-sulphobenzoate, prednisolone sodium 21-stearoglycolate, prednisolone tebutate, prednisolone 21-trimethylacetate, prednisone, prednival, prednylidene, prednylidene 21-diethylaminoacetate, tixocortol, triamcinolone, triamcinolone acetonide, triamcinolone benetonide and triamcinolone hexacetonide. Structurally related corticosteroids having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

In another specific embodiment, the present invention is directed to the use

of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment, or concurrent treatment) with any of one or more slow-acting antirheumatic drugs (SAARDs) or disease modifying antirheumatic drugs (DMARDS), prodrug esters, or pharmaceutically acceptable salts thereof for the treatment of the diseases and disorders recited herein, including acute and chronic inflammation such as rheumatic diseases, graft versus host disease, and multiple sclerosis. SAARDs or DMARDS, prodrug esters, and pharmaceutically acceptable salts thereof comprise: allocupreide sodium, auranofin, aurothioglucose, aurothioglycanide, azathioprine, brequinar sodium, bucillamine, calcium 3-aurothio-2-propanol-1-sulfonate, chlorambucil, chloroquine, clobuzarit, cuproxoline, cyclophosphamide, cyclosporin, dapsone, 15-deoxyspergualin, diacerein, glucosamine, gold salts (e.g., cycloquine gold salt, gold sodium thiomalate, gold sodium thiosulfate), hydroxychloroquine, hydroxychloroquine sulfate, hydroxyurea, kebuzone, levamisole, lobenzarit, melittin, 6-mercaptopurine, methotrexate, mizoribine, mycophenolate mofetil, myoral, nitrogen mustard, D-penicillamine, pyridinol imidazoles such as SKNF86002 and SB203580, rapamycin, thiols, thymopoietin and vincristine. Structurally related SAARDs or DMARDS having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

In another specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment, or concurrent treatment) with any of one or more COX2 inhibitors, prodrug esters, or pharmaceutically acceptable salts thereof for the treatment of the diseases and disorders recited herein, including acute and chronic inflammation. Examples of COX2 inhibitors, prodrug esters, or pharmaceutically acceptable salts thereof include, for example, celecoxib. Structurally related COX2 inhibitors having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

In still another specific embodiment, the present invention is directed to the use of an IL-1ra-L polypeptide in combination (pretreatment, post-treatment, or concurrent treatment) with any of one or more antimicrobials, prodrug esters,

or pharmaceutically acceptable salts thereof for the treatment of the diseases and disorders recited herein, including acute and chronic inflammation. Antimicrobials include, for example, the broad classes of penicillins, cephalosporins and other beta-lactams, aminoglycosides, azoles, quinolones, macrolides, rifamycins, tetracyclines, sulfonamides, lincosamides and polymyxins. The penicillins include, but are not limited to, penicillin G, penicillin V, methicillin, nafcillin, oxacillin, cloxacillin, dicloxacillin, floxacillin, ampicillin, ampicillin/sulbactam, amoxicillin, amoxicillin/clavulanate, hetacillin, cyclacillin, bacampicillin, carbenicillin, carbenicillin indanyl, ticarcillin, ticarcillin/clavulanate, azlocillin, mezlocillin, peperacillin, and mecillinam. The cephalosporins and other beta-lactams include, but are not limited to, cephalothin, cephapirin, cephalixin, cephradine, cefazolin, cefadroxil, cefaclor, cefamandole, cefotetan, cefoxitin, ceruroxime, cefonicid, ceforadine, cefixime, cefotaxime, moxalactam, ceftizoxime, ceftriaxone, cephaloperazone, ceftazidime, imipenem and aztreonam. The aminoglycosides include, but are not limited to, streptomycin, gentamicin, tobramycin, amikacin, netilmicin, kanamycin and neomycin. The azoles include, but are not limited to, fluconazole. The quinolones include, but are not limited to, nalidixic acid, norfloxacin, enoxacin, ciprofloxacin, ofloxacin, sparfloxacin and temafloxacin. The macrolides include, but are not limited to, erythromycin, spiramycin and azithromycin. The rifamycins include, but are not limited to, rifampin. The tetracyclines include, but are not limited to, spicycline, chlortetracycline, clomocycline, demeclocycline, deoxycycline, guamecycline, lymecycline, meclocycline, methacycline, minocycline, oxytetracycline, penimepicycline, pipacycline, rolitetracycline, sancycline, senociclin and tetracycline. The sulfonamides include, but are not limited to, sulfanilamide, sulfamethoxazole, sulfacetamide, sulfadiazine, sulfisoxazole and co-trimoxazole (trimethoprim/sulfamethoxazole). The lincosamides include, but are not limited to, clindamycin and lincomycin. The polymyxins (polypeptides) include, but are not limited to, polymyxin B and colistin.

Agonists or antagonists of IL-1RA-L polypeptide function may be used (simultaneously or sequentially) in combination with one or more cytokines,

growth factors, antibiotics, anti-inflammatories, and/or chemotherapeutic agents as is appropriate for the condition being treated.

Other diseases caused by or mediated by undesirable levels of one or more of IL-1, IL-1ra, or IL-1ra-L polypeptide are encompassed within the scope of the invention. Undesirable levels include excessive levels of IL-1, IL-1ra, or IL-1ra-L polypeptide and sub-normal levels of IL-1, IL-1ra, or IL-1ra-L polypeptide.

#### Uses of IL-1ra-L Nucleic Acids and Polypeptides

Nucleic acid molecules of the invention (including those that do not themselves encode biologically active polypeptides) may be used to map the locations of the IL-1ra-L gene and related genes on chromosomes. Mapping may be done by techniques known in the art, such as PCR amplification and *in situ* hybridization.

IL-1ra-L nucleic acid molecules (including those that do not themselves encode biologically active polypeptides), may be useful as hybridization probes in diagnostic assays to test, either qualitatively or quantitatively, for the presence of an IL-1ra-L nucleic acid molecule in mammalian tissue or bodily fluid samples.

Other methods may also be employed where it is desirable to inhibit the activity of one or more IL-1ra-L polypeptides. Such inhibition may be effected by nucleic acid molecules that are complementary to and hybridize to expression control sequences (triple helix formation) or to IL-1ra-L mRNA. For example, antisense DNA or RNA molecules, which have a sequence that is complementary to at least a portion of an IL-1ra-L gene can be introduced into the cell. Antisense probes may be designed by available techniques using the sequence of the IL-1ra-L gene disclosed herein. Typically, each such antisense molecule will be complementary to the start site (5' end) of each selected IL-1ra-L gene. When the antisense molecule then hybridizes to the corresponding IL-1ra-L mRNA, translation of this mRNA is prevented or reduced. Anti-sense inhibitors provide information relating to the decrease or absence of an IL-1ra-L polypeptide in a cell or organism.

Alternatively, gene therapy may be employed to create a dominant-

negative inhibitor of one or more IL-1ra-L polypeptides. In this situation, the DNA encoding a mutant polypeptide of each selected IL-1ra-L polypeptide can be prepared and introduced into the cells of a patient using either viral or non-viral methods as described herein. Each such mutant is typically designed to compete  
5 with endogenous polypeptide in its biological role.

In addition, an IL-1ra-L polypeptide, whether biologically active or not, may be used as an immunogen, that is, the polypeptide contains at least one epitope to which antibodies may be raised. Selective binding agents that bind to an IL-1ra-L polypeptide (as described herein) may be used for *in vivo* and *in vitro*  
10 diagnostic purposes, including, but not limited to, use in labeled form to detect the presence of IL-1ra-L polypeptide in a body fluid or cell sample. The antibodies may also be used to prevent, treat, or diagnose a number of diseases and disorders, including those recited herein. The antibodies may bind to an IL-1ra-L polypeptide so as to diminish or block at least one activity characteristic of an IL-  
15 1ra-L polypeptide, or may bind to a polypeptide to increase at least one activity characteristic of an IL-1ra-L polypeptide (including by increasing the pharmacokinetics of the IL-1ra-L polypeptide).

The IL-1ra-L polypeptides of the present invention can be used to clone IL-1ra-L polypeptide receptors, using an expression cloning strategy.  
20 Radiolabeled (<sup>125</sup>Iodine) IL-1ra-L polypeptide or affinity/activity-tagged IL-1ra-L polypeptide (such as an Fc fusion or an alkaline phosphatase fusion) can be used in binding assays to identify a cell type or cell line or tissue that expresses IL-1ra-L polypeptide receptors. RNA isolated from such cells or tissues can be converted to cDNA, cloned into a mammalian expression vector, and transfected  
25 into mammalian cells (such as COS or 293 cells) to create an expression library. A radiolabeled or tagged IL-1ra-L polypeptide can then be used as an affinity ligand to identify and isolate from this library the subset of cells that express the IL-1ra-L polypeptide receptors on their surface. DNA can then be isolated from these cells and transfected into mammalian cells to create a secondary expression  
30 library in which the fraction of cells expressing IL-1ra-L polypeptide receptors is many-fold higher than in the original library. This enrichment process can be

repeated iteratively until a single recombinant clone containing an IL-1ra-L polypeptide receptor is isolated. Isolation of the IL-1ra-L polypeptide receptors is useful for identifying or developing novel agonists and antagonists of the IL-1ra-L polypeptide signaling pathway. Such agonists and antagonists include soluble  
5 IL-1ra-L polypeptide receptors, anti-IL-1ra-L polypeptide receptor antibodies, small molecules, or antisense oligonucleotides, and they may be used for treating, preventing, or diagnosing one or more of the diseases or disorders described herein.

A deposit of cDNA encoding human IL-1ra-L polypeptide subcloned into  
10 pGEM-T easy and transfected into *E. coli* strain DH10B, having Accession No. PTA-1215, was made with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209 on January 20, 2000.

The following examples are intended for illustration purposes only, and should not be construed as limiting the scope of the invention in any way.

15

#### Example 1: Cloning of the Human IL-1ra-L Polypeptide Gene

Generally, materials and methods as described in Sambrook *et al. supra* were used to clone and analyze the gene encoding human IL-1ra-L polypeptide.

To isolate cDNA sequences encoding human IL-1ra-L polypeptide,  
20 homology-based BLAST searches of a human genomic database were performed. A 477 bp sequence (GA\_9383287\_422\_240) identified in this manner was found to share sequence homology with human IL-1ra. This sequence was used to design gene specific oligonucleotides for the identification of cDNA sources and the generation of cDNA clones, using various PCR strategies.

25 A number of cDNA libraries were analyzed in amplification reactions containing 10ng of cDNA library template DNA, 5 pmol each of the amplimers 2362-94 (5'-C-A-T-G-G-A-C-C-T-G-T-A-T-G-T-G-G-A-G-A-A-G-A-3'; SEQ ID NO: 9) and 2362-95 (5'-G-C-C-A-G-G-G-T-A-A-G-A-G-A-C-T-G-A-C-T-G-A-A-3'; SEQ ID NO: 10), and Ready-To-Go PCR beads (Amersham-Pharmacia,  
30 Piscataway, NJ) (Pharmacia, Piscataway, NJ), in a total reaction volume of 25  $\mu$ l. Reactions were performed at 95°C for 5 minutes for one cycle; 95°C for 15



seconds, 68°C for 15 seconds, and 72°C for 1 minute for 29 cycles; 72°C for 10 minutes for one cycle, and 95°C for 15 seconds, 68°C for 15 seconds, and 72°C for 1 minute for 10 cycles. A PCR product of the expected size (100 bp) was identified in a number of cDNA libraries, including libraries derived from a lymphoma cell line (oligo-dT and random primed), fetal kidney (oligo-dT and random primed), adult T-cells (oligo-dT primed), breast carcinoma, (oligo-dT and random primed), fetal lung (oligo-dT and random primed), fetal eye (oligo-dT and random primed), and fetal scalp (oligo-dT and random primed). The fetal scalp cDNA library was selected for further amplification experiments to isolate full-length cDNA sequences encoding IL-1ra-L polypeptide.

The fetal scalp cDNA library was prepared as follows. Total RNA was extracted from human fetal scalp using standard RNA extraction procedures and poly-A<sup>+</sup> RNA was selected from this total RNA using standard procedures. Random primed or oligo-dT primed cDNA was synthesized from this poly-A<sup>+</sup> RNA using the Superscript Plasmid System for cDNA Synthesis and Plasmid Cloning kit (Gibco-BRL), according to the manufacturer's suggested protocols, or other suitable procedure. The resulting cDNA was digested with appropriate restriction enzymes and was then ligated into pSPORT-1, or other suitable cloning vector. Ligation products were transformed into *E. coli* using standard techniques, and bacterial transformants were selected on culture plates containing either ampicillin, tetracycline, kanamycin, or chloramphenicol. The cDNA library consisted of all, or a subset, of these transformants.

Both 5'RACE and 3'RACE reactions were performed in order to generate the full-length cDNA sequence for IL-1ra-L polypeptide. To isolate cDNA sequences corresponding to the 5' end of the cDNA sequence for IL-1ra-L polypeptide, 5'RACE was performed using 10 ng of an random-primed human fetal scalp cDNA library in pSPORT1, 5 pmol each of the primers 870-02 (5'-A-G-C-G-G-A-T-A-A-C-A-A-T-T-T-C-A-C-A-C-A-G-G -3'; SEQ ID NO: 11) and 2366-21 (5'-G-C-C-T-A-G-G-C-T-G-G-A-T-T-T-A-T-T-C-C-A-C-A-G-3'; SEQ ID NO: 12), in a total reaction volume of 25 µl. Reactions were performed at 95°C for 1 minute for one cycle; holding at 80°C for the addition of 0.5 µl of

Advantage cDNA polymerase mix (Clontech); 95°C for 5 seconds, 64°C for 5 seconds, and 68°C for 3 minutes for 35 cycles; and 68°C for 3 minutes for one cycle. Nested PCR was performed using 10 µl of the 5'RACE amplification product (diluted 1/50) and the primers 1019-06 (5'-G-C-T-C-T-A-A-T-A-C-G-A-C-TC-A-C-T-A-T-A-G-G-G-3'; SEQ ID NO: 13) and 2362-98 (5'-C-T-G-A-T-G-T-G-G-T-G-G-A-GG-T-G-G-C-TA-T-3'; SEQ ID NO: 14), in a total reaction volume of 100 µl. Nested PCR Reactions were performed at 95°C for 1 minute for one cycle; holding at 80°C for the addition of 0.5 µl of Advantage cDNA polymerase mix; 95°C for 5 seconds, 64°C for 5 seconds, and 68°C for 3 minutes for 25 cycles; and 68°C for 3 minutes for one cycle. The amplification products obtained following nested PCR were isolated and sequenced directly.

To isolate cDNA sequences corresponding to the 3' end of the cDNA sequence for IL-1ra-L polypeptide, 3'RACE was performed using 10 ng of an oligo-dT-primed human fetal scalp cDNA library in pSPORT1 and the primers 1340-35 (5'-C-C-C-A-G-T-C-A-C-G-A-C-G-T-T-G-T-A-A-A-A-C-G-3'; SEQ ID NO: 15) and 2362-94, in a total reaction volume of 25 µl. Reactions were performed at 95°C for 1 minute for one cycle; holding at 80°C for the addition of 0.5 µl of Advantage cDNA polymerase mix; 95°C for 5 seconds, 64°C for 5 seconds, and 68°C for 3 minutes for 35 cycles; and 68°C for 3 minutes for one cycle. Nested PCR was performed using 10 µl of the 3'RACE amplification product (diluted 1/50) and the primers 1019-05 (5'-T-G-A-A-T-T-T-A-G-G-T-G-A-C-A-C-T-A-T-A-G-A-A-G-A-G-3'; SEQ ID NO: 16) and 2362-94, in a total reaction volume of 100 µl. Nested PCR Reactions were performed at 95°C for 1 minute for one cycle; holding at 80°C for the addition of 0.5 µl of Advantage cDNA polymerase mix (Clontech); 95°C for 5 seconds, 64°C for 5 seconds, and 68°C for 3 minutes for 25 cycles; and 68°C for 3 minutes for one cycle. The amplification products obtained following nested PCR were isolated and sequenced directly.

The sequences generated by 5' and 3'RACE form a contiguous sequence that appears to contain the full-length open reading frame for the IL-1ra-L gene. The full-length cDNA sequence for IL-1ra-L polypeptide was isolated by PCR

amplification using primers corresponding to the 5' and 3' ends of the IL-1ra-L gene as determined by 5' and 3'RACE. PCR was performed using 100 ng of an oligo-dT-primed human fetal scalp cDNA library and 10 pmoles each of the amplimers 2379-15 (5'-G-T-C-C-T-C-C-A-G-A-G-C-C-T-C-A-A-G-A-G-A-T-C-  
5 3'; SEQ ID NO: 17) and 2375-10 (5'-T-T-A-G-G-A-T-T-A-G-G-A-A-G-A-C-A-T-G-C-A-A-A-C-C-3'; SEQ ID NO: 18), in a total reaction volume of 50 µl. Reactions were performed at 95°C for 1 minute for one cycle; holding at 80°C for the addition of 1 µl of Advantage cDNA polymerase mix; 95°C for 5 seconds and 70°C for 3 minutes for 5 cycles; 95°C for 5 seconds, 68°C for 3 minutes for 5  
10 cycles; and 95°C for 5 seconds, 66°C for 5 seconds, and 66°C for 3 minutes for 25 cycles. The PCR product generated in this manner was isolated, cloned, and sequenced.

Sequence analysis of the predicted cDNA sequence for human IL-1ra-L polypeptide indicated that the gene comprises a 471 bp open reading frame  
15 encoding a protein of 157 amino acids (Figures 1A-1B). Figure 2 illustrates the amino acid sequence alignment of human IL-1δ (IL-1\_delta; SEQ ID NO: 3), human IL-1ra-L polypeptide (IL-1ra-L; SEQ ID NO: 2), human IL-1ε (IL-1\_epsilon; SEQ ID NO: 4), human IL-1 receptor antagonist, secreted polypeptide (IL-1ra\_sec; SEQ ID NO: 9), human IL-1β (IL-1\_beta; SEQ ID NO: 6), and a  
20 consensus sequence (consensus).

#### Example 2: IL-1ra-L mRNA Expression

PCR analysis was used to examine IL-1ra-L mRNA expression in various cDNA libraries. In one series of amplifications, reactions contained 10ng of  
25 cDNA library template DNA, 5 pmol of the amplimers 2362-94 and 2362-98, and Ready-To-Go PCR beads, in a total reaction volume of 25 µl. Reactions were performed at 95°C for 5 minutes for one cycle; 95°C for 15 seconds, 68°C for 15 seconds, and 72°C for 1 minute for 29 cycles; 72°C for 10 minutes for one cycle, and 95°C for 15 seconds, 68°C for 15 seconds, and 72°C for 1 minute for 10  
30 cycles. A PCR product of the expected size (100 bp) was identified in a number of cDNA libraries, including libraries derived from a lymphoma cell line (oligo-

dT and random primed), fetal kidney (oligo-dT and random primed), adult T-cells (oligo-dT primed), breast carcinoma, (olgio-dT and random primed), fetal lung (oligo-dT and random primed), fetal eye (oligo-dT and random primed), and fetal scalp (oligo-dT and random primed).

5        In another series of amplifications, reactions contained 10ng of Marathon™ cDNA (Clontech), 5 pmol of the amplimers 2362-94 and 2362-98, and 0.5 µl of Advantage cDNA polymerase mix, in a total reaction volume of 25 µl. Reactions were performed at 95°C for 1 minute for one cycle; 95°C for 5 seconds, 64°C for 5 seconds, and 68°C for 1 minute for 35 cycles; and 68°C for 1  
10       minute for one cycle. A PCR product of the expected size (100 bp) was identified in several cDNA sources, including human adult liver, lung, placenta, and spleen.

      The expression of IL-1ra-L mRNA can be examined by Northern blot analysis. Multiple human tissue northern blots (Clontech) are probed with a  
15       suitable restriction fragment isolated from a human IL-1ra-L polypeptide cDNA clone. The probe is labeled with <sup>32</sup>P-dCTP using standard techniques.

      Northern blots are prehybridized for 2 hours at 42°C in hybridization solution (5X SSC, 50% deionized formamide, 5X Denhardt's solution, 0.5% SDS, and 100 mg/ml denatured salmon sperm DNA) and then hybridized at 42°C  
20       overnight in fresh hybridization solution containing 5 ng/ml of the labeled probe. Following hybridization, the filters are washed twice for 10 minutes at room temperature in 2X SSC and 0.1% SDS, and then twice for 30 minutes at 65°C in 0.1X SSC and 0.1% SDS. The blots are then exposed to autoradiography.

25       The expression of IL-1ra-L mRNA is localized by *in situ* hybridization. A panel of normal embryonic and adult mouse tissues is fixed in 4% paraformaldehyde, embedded in paraffin, and sectioned at 5 µm. Sectioned tissues are permeabilized in 0.2 M HCl, digested with Proteinase K, and acetylated with triethanolamine and acetic anhydride. Sections are prehybridized  
30       for 1 hour at 60°C in hybridization solution (300 mM NaCl, 20 mM Tris-HCl, pH 8.0, 5 mM EDTA, 1X Denhardt's solution, 0.2% SDS, 10 mM DTT, 0.25 mg/ml

tRNA, 25 µg/ml polyA, 25 µg/ml polyC and 50% formamide) and then hybridized overnight at 60°C in the same solution containing 10% dextran and  $2 \times 10^4$  cpm/µl of a  $^{33}\text{P}$ -labeled antisense riboprobe complementary to the human IL-1ra-L gene. The riboprobe is obtained by *in vitro* transcription of a clone containing human  
5 IL-1ra-L cDNA sequences using standard techniques.

Following hybridization, sections are rinsed in hybridization solution, treated with RNaseA to digest unhybridized probe, and then washed in 0.1X SSC at 55°C for 30 minutes. Sections are then immersed in NTB-2 emulsion (Kodak, Rochester, NY), exposed for 3 weeks at 4°C, developed, and counterstained with  
10 hematoxylin and eosin. Tissue morphology and hybridization signal are simultaneously analyzed by darkfield and standard illumination for brain (one sagittal and two coronal sections), gastrointestinal tract (esophagus, stomach, duodenum, jejunum, ileum, proximal colon, and distal colon), pituitary, liver, lung, heart, spleen, thymus, lymph nodes, kidney, adrenal, bladder, pancreas,  
15 salivary gland, male and female reproductive organs (ovary, oviduct, and uterus in the female; and testis, epididymus, prostate, seminal vesicle, and vas deferens in the male), BAT and WAT (subcutaneous, peri-renal), bone (femur), skin, breast, and skeletal muscle.

### 20 Example 3: Production of IL-1ra-L Polypeptides

#### A. Expression of IL-1ra-L Polypeptides in Bacteria

PCR is used to amplify template DNA sequences encoding an IL-1ra-L polypeptide using primers corresponding to the 5' and 3' ends of the sequence. The amplified DNA products may be modified to contain restriction enzyme sites  
25 to allow for insertion into expression vectors. PCR products are gel purified and inserted into expression vectors using standard recombinant DNA methodology. An exemplary vector, such as pAMG21 (ATCC no. 98113) containing the lux promoter and a gene encoding kanamycin resistance is digested with Bam HI and Nde I for directional cloning of inserted DNA. The ligated mixture is transformed  
30 into an *E. coli* host strain by electroporation and transformants are selected for

kanamycin resistance. Plasmid DNA from selected colonies is isolated and subjected to DNA sequencing to confirm the presence of the insert.

Transformed host cells are incubated in 2xYT medium containing 30 µg/mL kanamycin at 30°C prior to induction. Gene expression is induced by the addition of N-(3-oxohexanoyl)-dl-homoserine lactone to a final concentration of 30 ng/mL followed by incubation at either 30°C or 37°C for six hours. The expression of IL-1ra-L polypeptide is evaluated by centrifugation of the culture, resuspension and lysis of the bacterial pellets, and analysis of host cell proteins by SDS-polyacrylamide gel electrophoresis.

Inclusion bodies containing IL-1ra-L polypeptide are purified as follows. Bacterial cells are pelleted by centrifugation and resuspended in water. The cell suspension is lysed by sonication and pelleted by centrifugation at 195,000 xg for 5 to 10 minutes. The supernatant is discarded, and the pellet is washed and transferred to a homogenizer. The pellet is homogenized in 5 mL of a Percoll solution (75% liquid Percoll and 0.15 M NaCl) until uniformly suspended and then diluted and centrifuged at 21,600 xg for 30 minutes. Gradient fractions containing the inclusion bodies are recovered and pooled. The isolated inclusion bodies are analyzed by SDS-PAGE.

A single band on an SDS polyacrylamide gel corresponding to *E. coli*-produced IL-1ra-L polypeptide is excised from the gel, and the N-terminal amino acid sequence is determined essentially as described by Matsudaira *et al.*, 1987, *J. Biol. Chem.* 262:10-35.

#### B. Expression of IL-1ra-L Polypeptide in Mammalian Cells

PCR is used to amplify template DNA sequences encoding an IL-1ra-L polypeptide using primers corresponding to the 5' and 3' ends of the sequence. The amplified DNA products may be modified to contain restriction enzyme sites to allow for insertion into expression vectors. PCR products are gel purified and inserted into expression vectors using standard recombinant DNA methodology.

An exemplary expression vector, pCEP4 (Invitrogen, Carlsbad, CA), that contains an Epstein-Barr virus origin of replication, may be used for the expression of IL-

1ra-L polypeptides in 293-EBNA-1 cells. Amplified and gel purified PCR products are ligated into pCEP4 vector and introduced into 293-EBNA cells by lipofection. The transfected cells are selected in 100 µg/mL hygromycin and the resulting drug-resistant cultures are grown to confluence. The cells are then  
5 cultured in serum-free media for 72 hours. The conditioned media is removed and IL-1ra-L polypeptide expression is analyzed by SDS-PAGE.

IL-1ra-L polypeptide expression may be detected by silver staining. Alternatively, IL-1ra-L polypeptide is produced as a fusion protein with an epitope tag, such as an IgG constant domain or a FLAG epitope, which may be  
10 detected by Western blot analysis using antibodies to the peptide tag.

IL-1ra-L polypeptides may be excised from an SDS-polyacrylamide gel, or IL-1ra-L fusion proteins are purified by affinity chromatography to the epitope tag, and subjected to N-terminal amino acid sequence analysis as described  
15 herein.

### C. Expression and Purification of IL-1ra-L Polypeptide in Mammalian Cells

IL-1ra-L polypeptide expression constructs are introduced into 293 EBNA or CHO cells using either a lipofection or calcium phosphate protocol.

To conduct functional studies on the IL-1ra-L polypeptides that are  
20 produced, large quantities of conditioned media are generated from a pool of hygromycin selected 293 EBNA clones. The cells are cultured in 500 cm Nunc Triple Flasks to 80% confluence before switching to serum free media a week prior to harvesting the media. Conditioned media is harvested and frozen at -20°C until purification.

25 Conditioned media is purified by affinity chromatography as described below. The media is thawed and then passed through a 0.2 µm filter. A Protein G column is equilibrated with PBS at pH 7.0, and then loaded with the filtered media. The column is washed with PBS until the absorbance at A<sub>280</sub> reaches a baseline. IL-1ra-L polypeptide is eluted from the column with 0.1 M Glycine-HCl at pH 2.7 and immediately neutralized with 1 M Tris-HCl at pH 8.5.  
30

Fractions containing IL-1ra-L polypeptide are pooled, dialyzed in PBS, and stored at -70°C.

For Factor Xa cleavage of the human IL-1ra-L polypeptide-Fc fusion polypeptide, affinity chromatography-purified protein is dialyzed in 50 mM Tris-HCl, 100 mM NaCl, 2 mM CaCl<sub>2</sub> at pH 8.0. The restriction protease Factor Xa is added to the dialyzed protein at 1/100 (w/w) and the sample digested overnight at room temperature.

#### Example 4: Production of Anti-IL-1ra-L Polypeptide Antibodies

10 Antibodies to IL-1ra-L polypeptides may be obtained by immunization with purified protein or with IL-1ra-L peptides produced by biological or chemical synthesis. Suitable procedures for generating antibodies include those described in Hudson and Bay, *Practical Immunology* (2nd ed., Blackwell Scientific Publications).

15 In one procedure for the production of antibodies, animals (typically mice or rabbits) are injected with an IL-1ra-L antigen (such as an IL-1ra-L polypeptide), and those with sufficient serum titer levels as determined by ELISA are selected for hybridoma production. Spleens of immunized animals are collected and prepared as single cell suspensions from which splenocytes are recovered. The splenocytes are fused to mouse myeloma cells (such as Sp2/0-Ag14 cells), are first incubated in DMEM with 200 U/mL penicillin, 200 µg/mL streptomycin sulfate, and 4 mM glutamine, and are then incubated in HAT selection medium (hypoxanthine, aminopterin, and thymidine). After selection, the tissue culture supernatants are taken from each fusion well and tested for anti-IL-1ra-L antibody production by ELISA.

25 Alternative procedures for obtaining anti-IL-1ra-L antibodies may also be employed, such as the immunization of transgenic mice harboring human Ig loci for production of human antibodies, and the screening of synthetic antibody libraries, such as those generated by mutagenesis of an antibody variable domain.

30

#### Example 5: Expression of IL-1ra-L Polypeptide in Transgenic Mice



To assess the biological activity of IL-1ra-L polypeptide, a construct encoding an IL-1ra-L polypeptide/Fc fusion protein under the control of a liver specific ApoE promoter is prepared. The delivery of this construct is expected to cause pathological changes that are informative as to the function of IL-1ra-L polypeptide. Similarly, a construct containing the full-length IL-1ra-L polypeptide under the control of the beta actin promoter is prepared. The delivery of this construct is expected to result in ubiquitous expression.

To generate these constructs, PCR is used to amplify template DNA sequences encoding an IL-1ra-L polypeptide using primers that correspond to the 5' and 3' ends of the desired sequence and which incorporate restriction enzyme sites to permit insertion of the amplified product into an expression vector. Following amplification, PCR products are gel purified, digested with the appropriate restriction enzymes, and ligated into an expression vector using standard recombinant DNA techniques. For example, amplified IL-1ra-L polypeptide sequences can be cloned into an expression vector under the control of the human  $\beta$ -actin promoter as described by Graham *et al.*, 1997, *Nature Genetics*, 17:272-74 and Ray *et al.*, 1991, *Genes Dev.* 5:2265-73.

Following ligation, reaction mixtures are used to transform an *E. coli* host strain by electroporation and transformants are selected for drug resistance. Plasmid DNA from selected colonies is isolated and subjected to DNA sequencing to confirm the presence of an appropriate insert and absence of mutation. The IL-1ra-L polypeptide expression vector is purified through two rounds of CsCl density gradient centrifugation, cleaved with a suitable restriction enzyme, and the linearized fragment containing the IL-1ra-L polypeptide transgene is purified by gel electrophoresis. The purified fragment is resuspended in 5 mM Tris, pH 7.4, and 0.2 mM EDTA at a concentration of 2 mg/mL.

Single-cell embryos from BDF1 x BDF1 bred mice are injected as described (PCT Pub. No. WO 97/23614). Embryos are cultured overnight in a CO<sub>2</sub> incubator and 15-20 two-cell embryos are transferred to the oviducts of a pseudopregnant CD1 female mice. Offspring obtained from the implantation of microinjected embryos are screened by PCR amplification of the integrated

transgene in genomic DNA samples as follows. Ear pieces are digested in 20 mL ear buffer (20 mM Tris, pH 8.0, 10 mM EDTA, 0.5% SDS, and 500 mg/mL proteinase K) at 55°C overnight. The sample is then diluted with 200 mL of TE, and 2 mL of the ear sample is used in a PCR reaction using appropriate primers.

5           At 8 weeks of age, transgenic founder animals and control animals are sacrificed for necropsy and pathological analysis. Portions of spleen are removed and total cellular RNA isolated from the spleens using the Total RNA Extraction Kit (Qiagen) and transgene expression determined by RT-PCR. RNA recovered from spleens is converted to cDNA using the SuperScript™ Preamplification  
10   System (Gibco-BRL) as follows. A suitable primer, located in the expression vector sequence and 3' to the IL-1ra-L polypeptide transgene, is used to prime cDNA synthesis from the transgene transcripts. Ten mg of total spleen RNA from transgenic founders and controls is incubated with 1 mM of primer for 10 minutes at 70°C and placed on ice. The reaction is then supplemented with 10 mM Tris-  
15   HCl, pH 8.3, 50 mM KCl, 2.5 mM MgCl<sub>2</sub>, 10 mM of each dNTP, 0.1 mM DTT, and 200 U of SuperScript II reverse transcriptase. Following incubation for 50 minutes at 42°C, the reaction is stopped by heating for 15 minutes at 72°C and digested with 2U of RNase H for 20 minutes at 37°C. Samples are then amplified by PCR using primers specific for IL-1ra-L polypeptide.

20

#### Example 6: Biological Activity of IL-1ra-L Polypeptide in Transgenic Mice

Prior to euthanasia, transgenic animals are weighed, anesthetized by isofluorane and blood drawn by cardiac puncture. The samples are subjected to hematology and serum chemistry analysis. Radiography is performed after  
25   terminal exsanguination. Upon gross dissection, major visceral organs are subject to weight analysis.

Following gross dissection, tissues (*i.e.*, liver, spleen, pancreas, stomach, the entire gastrointestinal tract, kidney, reproductive organs, skin and mammary glands, bone, brain, heart, lung, thymus, trachea, esophagus, thyroid, adrenals,  
30   urinary bladder, lymph nodes and skeletal muscle) are removed and fixed in 10% buffered Zn-Formalin for histological examination. After fixation, the tissues are

processed into paraffin blocks, and 3 mm sections are obtained. All sections are stained with hematoxylin and eosin, and are then subjected to histological analysis.

5 The spleen, lymph node, and Peyer's patches of both the transgenic and the control mice are subjected to immunohistology analysis with B cell and T cell specific antibodies as follows. The formalin fixed paraffin embedded sections are deparaffinized and hydrated in deionized water. The sections are quenched with 3% hydrogen peroxide, blocked with Protein Block (Lipshaw, Pittsburgh, PA), and incubated in rat monoclonal anti-mouse B220 and CD3 (Harlan, Indianapolis, IN). Antibody binding is detected by biotinylated rabbit anti-rat immunoglobulins and peroxidase conjugated streptavidin (BioGenex, San Ramon, CA) with DAB as a chromagen (BioTek, Santa Barbara, CA). Sections are counterstained with hematoxylin.

15 After necropsy, MLN and sections of spleen and thymus from transgenic animals and control littermates are removed. Single cell suspensions are prepared by gently grinding the tissues with the flat end of a syringe against the bottom of a 100 mm nylon cell strainer (Becton Dickinson, Franklin Lakes, NJ). Cells are washed twice, counted, and approximately  $1 \times 10^6$  cells from each tissue are then incubated for 10 minutes with 0.5  $\mu$ g CD16/32(Fc $\gamma$ III/II) Fc block in a 20  $\mu$ L volume. Samples are then stained for 30 minutes at 2-8°C in a 100  $\mu$ L volume of PBS (lacking Ca<sup>+</sup> and Mg<sup>+</sup>), 0.1% bovine serum albumin, and 0.01% sodium azide with 0.5  $\mu$ g antibody of FITC or PE-conjugated monoclonal antibodies against CD90.2 (Thy-1.2), CD45R (B220), CD11b(Mac-1), Gr-1, CD4, or CD8 (PharMingen, San Diego, CA). Following antibody binding, the cells are washed and then analyzed by flow cytometry on a FACScan (Becton Dickinson).

25 While the present invention has been described in terms of the preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations that come within the scope of the invention as claimed.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- 5 (a) the nucleotide sequence as set forth in SEQ ID NO: 1;
- (b) the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-1215;
- (c) a nucleotide sequence encoding the polypeptide as set forth in SEQ ID NO: 2;
- 10 (d) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (c); and
- (e) a nucleotide sequence complementary to any of (a) - (c).

2. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- 15 (a) a nucleotide sequence encoding a polypeptide which is at least about 70 percent identical to the polypeptide as set forth in SEQ ID NO: 2, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;
- 20 (b) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in SEQ ID NO: 1, the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-1215, or (a);
- (c) a region of the nucleotide sequence of SEQ ID NO: 1, the DNA insert in ATCC Deposit No. PTA-1215, (a), or (b) encoding a polypeptide
- 25 fragment of at least about 25 amino acid residues, wherein the polypeptide fragment has an activity of the encoded polypeptide as set forth in SEQ ID NO: 2, or is antigenic;
- (d) a region of the nucleotide sequence of SEQ ID NO: 1, the DNA insert in ATCC Deposit No. PTA-1215, or any of (a) - (c) comprising a fragment
- 30 of at least about 16 nucleotides;

(e) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (d); and

(f) a nucleotide sequence complementary to any of (a) - (d).

5           3. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one conservative amino acid substitution, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

10           (b) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid insertion, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(c) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid deletion, wherein the encoded polypeptide  
15 has an activity of the polypeptide set forth in SEQ ID NO: 2;

(d) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 which has a C- and/or N- terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(e) a nucleotide sequence encoding a polypeptide as set forth in SEQ  
20 ID NO: 2 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(f) a nucleotide sequence of any of (a) - (e) comprising a fragment of  
25 at least about 16 nucleotides;

(g) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (f); and

(h) a nucleotide sequence complementary to any of (a) - (e).

30           4. A vector comprising the nucleic acid molecule of any of Claims 1, 2, or 3.

5. A host cell comprising the vector of Claim 4.

6. The host cell of Claim 5 that is a eukaryotic cell.

5

7. The host cell of Claim 5 that is a prokaryotic cell.

8. A process of producing an IL-1ra-L polypeptide comprising culturing the host cell of Claim 5 under suitable conditions to express the polypeptide, and optionally isolating the polypeptide from the culture.

10

9. A polypeptide produced by the process of Claim 8.

10. The process of Claim 8, wherein the nucleic acid molecule comprises promoter DNA other than the promoter DNA for the native IL-1ra-L polypeptide operatively linked to the DNA encoding the IL-1ra-L polypeptide.

15

11. The isolated nucleic acid molecule according to Claim 2, wherein the percent identity is determined using a computer program selected from the group consisting of GAP, BLASTN, FASTA, BLASTA, BLASTX, BestFit, and the Smith-Waterman algorithm.

20

12. A process for determining whether a compound inhibits IL-1ra-L polypeptide activity or IL-1ra-L polypeptide production comprising exposing a cell according to any of Claims 5, 6, or 7 to the compound and measuring IL-1ra-L polypeptide activity or IL-1ra-L polypeptide production in said cell.

25

13. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

30

(a) the amino acid sequence as set forth in SEQ ID NO: 2; and

(b) the amino acid sequence encoded by the DNA insert in ATCC Deposit No. PTA-1215.

14. An isolated polypeptide comprising the amino acid sequence  
5 selected from the group consisting of:

- (a) an amino acid sequence for an ortholog of SEQ ID NO: 2;
- (b) an amino acid sequence which is at least about 70 percent identical to the amino acid sequence of SEQ ID NO: 2, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;
- 10 (c) a fragment of the amino acid sequence set forth in SEQ ID NO: 2 comprising at least about 25 amino acid residues, wherein the fragment has an activity of the polypeptide set forth in SEQ ID NO: 2, or is antigenic; and
- (d) an amino acid sequence for an allelic variant or splice variant of the amino acid sequence as set forth in SEQ ID NO: 2, the amino acid sequence  
15 encoded by the DNA insert in ATCC Deposit No. PTA-1215, (a), or (b).

15. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence as set forth in SEQ ID NO: 2 with at least  
20 one conservative amino acid substitution, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;
- (b) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one amino acid insertion, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;
- 25 (c) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one amino acid deletion, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;
- (d) the amino acid sequence as set forth in SEQ ID NO: 2 which has a C- and/or N- terminal truncation, wherein the polypeptide has an activity of the  
30 polypeptide set forth in SEQ ID NO: 2; and

(e) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2.

16. An isolated polypeptide encoded by the nucleic acid molecule of any of Claims 1, 2, or 3, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2.

10

17. The isolated polypeptide according to Claim 14, wherein the percent identity is determined using a computer program selected from the group consisting of GAP, BLASTP, FASTA, BLASTA, BLASTX, BestFit, and the Smith-Waterman algorithm.

15

18. A selective binding agent or fragment thereof which specifically binds the polypeptide of any of Claims 13, 14, or 15.

19. The selective binding agent or fragment thereof of Claim 18 that specifically binds the polypeptide comprising the amino acid sequence as set forth in SEQ ID NO: 2, or a fragment thereof.

20

20. The selective binding agent of Claim 18 that is an antibody or fragment thereof.

25

21. The selective binding agent of Claim 18 that is a humanized antibody.

22. The selective binding agent of Claim 18 that is a human antibody or fragment thereof.

30



23. The selective binding agent of Claim 18 that is a polyclonal antibody or fragment thereof.

24. The selective binding agent Claim 18 that is a monoclonal antibody or fragment thereof.

25. The selective binding agent of Claim 18 that is a chimeric antibody or fragment thereof.

26. The selective binding agent of Claim 18 that is a CDR-grafted antibody or fragment thereof.

27. The selective binding agent of Claim 18 that is an antiidiotypic antibody or fragment thereof.

28. The selective binding agent of Claim 18 that is a variable region fragment.

29. The variable region fragment of Claim 28 that is a Fab or a Fab' fragment.

30. A selective binding agent or fragment thereof comprising at least one complementarity determining region with specificity for a polypeptide having the amino acid sequence of SEQ ID NO: 2.

31. The selective binding agent of Claim 18 that is bound to a detectable label.

32. The selective binding agent of Claim 18 that antagonizes IL-1ra-L polypeptide biological activity.

33. A method for treating, preventing, or ameliorating an IL-1ra-L polypeptide-related disease, condition, or disorder comprising administering to a patient an effective amount of a selective binding agent according to Claim 18.

5 34. A selective binding agent produced by immunizing an animal with a polypeptide comprising an amino acid sequence of SEQ ID NO: 2.

35. A hybridoma which produces a selective binding agent which is capable of binding a polypeptide according to any of Claims 1, 2, or 3.

10

36. A method of detecting or quantitating the amount of IL-1ra-L polypeptide using the anti-IL-1ra-L antibody or fragment of Claim 18.

37. A composition comprising the polypeptide of any of Claims 13, 14, or 15, and a pharmaceutically acceptable formulation agent.

15

38. The composition of Claim 37, wherein the pharmaceutically acceptable formulation agent is a carrier, adjuvant, solubilizer, stabilizer, or anti-oxidant.

20

39. A polypeptide comprising a derivative of the polypeptide of any of Claims 13, 14, or 15.

40. The polypeptide of Claim 39 that is covalently modified with a water-soluble polymer.

25

41. The polypeptide of Claim 40, wherein the water-soluble polymer is selected from the group consisting of polyethylene glycol, monomethoxy-polyethylene glycol, dextran, cellulose, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols, and polyvinyl alcohol.

30

42. A composition comprising a nucleic acid molecule of any of Claims 1, 2, or 3 and a pharmaceutically acceptable formulation agent.

5 43. The composition of Claim 42, wherein said nucleic acid molecule is contained in a viral vector.

44. A viral vector comprising a nucleic acid molecule of any of Claims 1, 2, or 3.

10

45. A fusion polypeptide comprising the polypeptide of any of Claims 13, 14, or 15 fused to a heterologous amino acid sequence.

46. The fusion polypeptide of Claim 45, wherein the heterologous amino acid sequence is an IgG constant domain or fragment thereof.

15

47. A method for treating, preventing, or ameliorating a medical condition comprising administering to a patient the polypeptide of any of Claims 13, 14, or 15, or the polypeptide encoded by the nucleic acid of any of Claims 1, 2, or 3.

20

48. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

(a) determining the presence or amount of expression of the polypeptide of any of Claims 13, 14, or 15, or the polypeptide encoded by the nucleic acid molecule of any of Claims 1, 2, or 3 in a sample; and

25

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.

30

49. A device, comprising:

- (a) a membrane suitable for implantation; and
  - (b) cells encapsulated within said membrane, wherein said cells secrete a protein of any of Claims 13, 14, or 15; and
- said membrane is permeable to said protein and impermeable to materials  
5 detrimental to said cells.

50. A method of identifying a compound which binds to an IL-1ra-L polypeptide comprising:

- (a) contacting the polypeptide of any of Claims 13, 14, or 15 with a  
10 compound; and
- (b) determining the extent of binding of the IL-1ra-L polypeptide to the compound.

51. The method of Claim 50, further comprising determining the  
15 activity of the polypeptide when bound to the compound.

52. A method of modulating levels of a polypeptide in an animal comprising administering to the animal the nucleic acid molecule of any of Claims 1, 2, or 3.

20

53. A transgenic non-human mammal comprising the nucleic acid molecule of any of Claims 1, 2, or 3.

54. A process for determining whether a compound inhibits IL-1ra-L  
25 polypeptide activity or IL-1ra-L polypeptide production comprising exposing a transgenic mammal according to Claim 53 to the compound, and measuring IL-1ra-L polypeptide activity or IL-1ra-L polypeptide production in said mammal.

The present invention provides novel Interleukin-1 Receptor Antagonist-Like (IL-1ra-L) polypeptides and nucleic acid molecules encoding the same. The invention also provides selective binding agents, vectors, host cells, and methods for producing IL-1ra-L polypeptides. The invention further provides pharmaceutical compositions and methods for the diagnosis, treatment, amelioration, and/or prevention of diseases, disorders, and conditions associated with IL-1ra-L polypeptides.

FIG. 1A

```

gtgttgctcc actgtcagtc ctccagagcc tcaagagatc tttgggcat atcagctttc 60
tttccaaaat gaacacaccc aggggcagga aagaatgctc tttccttggt cattaagggg 120
cctgggagtc ctggaccagc ttttcatgca gctagaccac ttacatgcaa ctagagcctt 180
gactttgaaa cgagggacaa aagcatctct tgctaaaggt aactttctgct gcttagaacc 240
cagcctcttc accaccatct gatctatctt gttctcttca caaaaggctc tgaagacatc 300

atg aac cca caa cgg gag gca gca ccc aaa tcc tat gct att cgt gat 348
Met Asn Pro Gln Arg Glu Ala Ala Pro Lys Ser Tyr Ala Ile Arg Asp
  1           5           10          15

tct cga cag atg gtg tgg gtc ctg agt gga aat tct tta ata gca gct 396
Ser Arg Gln Met Val Trp Val Leu Ser Gly Asn Ser Leu Ile Ala Ala
          20           25           30

cct ctt agc cgc agc att aag cct gtc act ctt cat tta ata gcc tgt 444
Pro Leu Ser Arg Ser Ile Lys Pro Val Thr Leu His Leu Ile Ala Cys
          35           40           45

aga gac aca gaa ttc agt gac aag gaa aag ggt aat atg gtt tac ctg 492
Arg Asp Thr Glu Phe Ser Asp Lys Glu Lys Gly Asn Met Val Tyr Leu
          50           55           60

gga atc aag gga aaa gat ctc tgt ctc ttc tgt gca gaa att cag ggc 540
Gly Ile Lys Gly Lys Asp Leu Cys Leu Phe Cys Ala Glu Ile Gln Gly
          65           70           75           80

aag cct act ttg cag ctt aag gaa aaa aat atc atg gac ctg tat gtg 588
Lys Pro Thr Leu Gln Leu Lys Glu Lys Asn Ile Met Asp Leu Tyr Val
          85           90           95

gag aag aaa gca cag aag ccc ttt ctc ttt ttc cac aat aaa gaa ggc 636
Glu Lys Lys Ala Gln Lys Pro Phe Leu Phe Phe His Asn Lys Glu Gly
          100          105          110

tcc act tct gtc ttt cag tca gtc tct tac cct ggc tgg ttc ata gcc 684
Ser Thr Ser Val Phe Gln Ser Val Ser Tyr Pro Gly Trp Phe Ile Ala
          115          120          125

acc tcc acc aca tca gga cag ccc atc ttt ctc acc aag gag aga ggc 732
Thr Ser Thr Thr Ser Gly Gln Pro Ile Phe Leu Thr Lys Glu Arg Gly
          130          135          140

ata act aat aac act aac ttc tac tta gat tct gtg gaa taa 774
Ile Thr Asn Asn Thr Asn Phe Tyr Leu Asp Ser Val Glu
          145          150          155

atccagccta ggctgtgggt ggctggttcc aggatagaga atcaagctgt cagagtcac 834
ttaacagatc attatgcgac tgagttcact agcagttcag cccatccata gcttacctca 894

```

GenBank accession number: U00096

1

ttcttactat	ccaaaagcca	cctcctcctc	caaacatcca	tttctgtacc	aagaccctca	954
ctcgaatgtc	actatcccaa	gatgaaacct	aaaaatcact	ttccattctt	tcttgatctt	1014
accccaccat	ccactcagct	gccatgcca	gtttagttaa	ccccccaaat	gctgcttcat	1074
gcaaccttcc	attcctattc	cttttgccaa	cccatgatgt	agagatgtgg	attcatgaca	1134
ttttgttcat	acaacttctt	caataaaaaca	ttataatatg	tgccccaaag	ataaagctga	1194
agaatgagat	gaatgtgaaa	ttaaagggtt	gcatgtcttc	ctaatacctaa		1244

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
2	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36	38	40	42	44	46	48	50	52	54	56	58	60	62	64	66	68	70	72	74	76	78	80	82	84	86	88	90	92	94	96	98	100	102	104	106	108	110	112	114	116	118	120	122	124	126	128	130	132	134	136	138	140	142	144	146	148	150	152	154	156	158	160	162	164	166	168	170	172	174	176	178	180	182	184	186	188	190	192	194	196	198	200
3	3	6	9	12	15	18	21	24	27	30	33	36	39	42	45	48	51	54	57	60	63	66	69	72	75	78	81	84	87	90	93	96	99	102	105	108	111	114	117	120	123	126	129	132	135	138	141	144	147	150	153	156	159	162	165	168	171	174	177	180	183	186	189	192	195	198	201	204	207	210	213	216	219	222	225	228	231	234	237	240	243	246	249	252	255	258	261	264	267	270	273	276	279	282	285	288	291	294	297	300
4	4	8	12	16	20	24	28	32	36	40	44	48	52	56	60	64	68	72	76	80	84	88	92	96	100	104	108	112	116	120	124	128	132	136	140	144	148	152	156	160	164	168	172	176	180	184	188	192	196	200	204	208	212	216	220	224	228	232	236	240	244	248	252	256	260	264	268	272	276	280	284	288	292	296	300																									
5	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95	100	105	110	115	120	125	130	135	140	145	150	155	160	165	170	175	180	185	190	195	200	205	210	215	220	225	230	235	240	245	250	255	260	265	270	275	280	285	290	295	300																																								
6	6	12	18	24	30	36	42	48	54	60	6																																																																																									

FIG. 2

	1				50
IL-1_delta	~~~~~	~~~~~	~~~~~MNPQ	REAAPKSYAI	RDSRQMVWVL
IL-1ra-L	~~~~~	~~~~~	~~~~~MNPQ	REAAPKSYAI	RDSRQMVWVL
IL-1_epsilon	~~~~~	~~~~~	~~~~~MEKAL	KIDTPQQGSI	QDINHRVWVL
IL-1ra_sec	MEICRGLRSH	LITLLLFLFH	SETICRPSGR	KSSKMQAFRI	WDVNQKTFYL
IL-1_beta	~~~~~	~~~~~	~~~~~	APVRSINCTL	RDSQQKSLVM
consensus	-----	-----	-----MNPQ	-EAAP-SYAI	RDS-Q-VWVL
	51				100
IL-1_delta	SG.NSLIAAP	LSRSIKPVTI	HLIACRDTEF	SDKEKGNMVY	LGIKGDLC
IL-1ra-L	SG.NSLIAAP	LSRSIKPVTI	HLIACRDTEF	SDKEKGNMVY	LGIKGDLC
IL-1_epsilon	QD.QTLIAVP	RKDRMSPVTI	ALISCRHVET	LEKDRGNPIY	LGLNGLNLCL
IL-1ra_sec	RN.NQLVAGY	LQ...GP.NV	NLEEKIDVVP	IEP...HALF	LGIHGGKMCL
IL-1_beta	SGPYELKALH	LQGQDMEQQV	.VFSMSFVQG	EESNDKIPVA	LGLKEKNLYL
consensus	SG-NSLIAAP	L-RSIKPVT-	HLI-CRDVEF	SEKEKGN-VY	LGIKGL-LCL
	101				150
IL-1_delta	FCAEIQ GKPT	LQLKLQGSQD	.NIGKDT CWK	LVGIHTCINL	DVRESCFMG.
IL-1ra-L	FCAEIQ GKPT	LQLKEKNIMD	LYVEKKAQKP	FLFFH...NK	EGSTSVFQSV
IL-1_epsilon	MCAKVG DQPT	LQLKEKDIMD	LYNQPEPVKS	FLFYH...SQ	SGRNSTFESV
IL-1ra_sec	SCVKSGDETR	LQLEAVNITD	LSENRKQDKR	FAFIR...SD	SGPTTSFESA
IL-1_beta	SCVLKDDKPT	LQLESVDPKN	.YPKKKMEKR	FVFNKIEINN	KLE...FESA
consensus	-CA-I-DKPT	LQLKE--IMD	LY--KK--KR	F-FIH--IN-	SGRTS-FES-
	151				193
IL-1_delta	TLDQWGIGVG	RKKWKSSFQH	HHLRKKDKDF	SSMRTNIGMP	GRM
IL-1ra-L	SYPGWFIATS	TTSGQPIF..	...LTKERGI	TN.NTNFYLD	SVE
IL-1_epsilon	AFPGWFIAMS	SEGGCPLILT	QELGKANTTD	FGLTMLF~~~	~~~
IL-1ra_sec	ACPGWFLCTA	MEADQPVSLT	N...MPDEGV	MVTKFYFQED	E~~
IL-1_beta	QFPNWIYSTS	QAENMPVFLG	GTKGGQDITD	FTMQFVSS~~	~~~
consensus	AFPGWFIATS	-E-GQPVFLT	--LGKKD--D	F-M--NF--D	---



# SEQUENCE LISTING

<110> Welcher, Andrew A.  
Jing, Shugian  
Luethy, Roland

<120> Interleukin-1 Receptor Antagonist-Like Molecules and  
Uses Thereof

<130> 00-1214

<140>

<141>

<150> 60/170,052

<151> 1999-12-10

<160> 18

<170> PatentIn Ver. 2.0

<210> 1

<211> 1244

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (301)..(774)

<400> 1

```
gtgttgctcc actgtcagtc ctccagagcc tcaagagatc tttgggccat atcagctttc 60
tttccaaaat gaacacaccc aggggcagga aagaatgctc tttccttggt cattaagggg 120
cctggggagtc ctggaccagc ttttcatgca gctagaccac ttacatgcaa ctagagcctt 180
gactttgaaa cgaggggacaa aagcatctct tgctaaaggt aacttctgct gcttagaacc 240
cagcctcctc accaccatct gatctatctt gttctcttca caaaaggctc tgaagacatc 300

atg aac cca caa cgg gag gca gca ccc aaa tcc tat gct att cgt gat 348
Met Asn Pro Gln Arg Glu Ala Ala Pro Lys Ser Tyr Ala Ile Arg Asp
  1           5           10          15

tct cga cag atg gtg tgg gtc ctg agt gga aat tct tta ata gca gct 396
Ser Arg Gln Met Val Trp Val Leu Ser Gly Asn Ser Leu Ile Ala Ala
      20           25           30

cct ctt agc cgc agc att aag cct gtc act ctt cat tta ata gcc tgt 444
Pro Leu Ser Arg Ser Ile Lys Pro Val Thr Leu His Leu Ile Ala Cys
      35           40           45

aga gac aca gaa ttc agt gac aag gaa aag ggt aat atg gtt tac ctg 492
Arg Asp Thr Glu Phe Ser Asp Lys Glu Lys Gly Asn Met Val Tyr Leu
      50           55           60
```

gga atc aag gga aaa gat ctc tgt ctc ttc tgt gca gaa att cag ggc 540  
 Gly Ile Lys Gly Lys Asp Leu Cys Leu Phe Cys Ala Glu Ile Gln Gly  
 65 70 75 80

aag cct act ttg cag ctt aag gaa aaa aat atc atg gac ctg tat gtg 588  
 Lys Pro Thr Leu Gln Leu Lys Glu Lys Asn Ile Met Asp Leu Tyr Val  
 85 90 95

gag aag aaa gca cag aag ccc ttt ctc ttt ttc cac aat aaa gaa ggc 636  
 Glu Lys Lys Ala Gln Lys Pro Phe Leu Phe Phe His Asn Lys Glu Gly  
 100 105 110

tcc act tct gtc ttt cag tca gtc tct tac cct ggc tgg ttc ata gcc 684  
 Ser Thr Ser Val Phe Gln Ser Val Ser Tyr Pro Gly Trp Phe Ile Ala  
 115 120 125

acc tcc acc aca tca gga cag ccc atc ttt ctc acc aag gag aga ggc 732  
 Thr Ser Thr Thr Ser Gly Gln Pro Ile Phe Leu Thr Lys Glu Arg Gly  
 130 135 140

ata act aat aac act aac ttc tac tta gat tct gtg gaa taa 774  
 Ile Thr Asn Asn Thr Asn Phe Tyr Leu Asp Ser Val Glu  
 145 150 155

atccagccta ggctgtgggt ggctggttcc aggatagaga atcaagctgt cagagtcac 834  
 ttaacagatc attatgcgac tgagttcact agcagttcag cccatccata gcttacctca 894  
 ttcttactat ccaaaagcca cctcctcctc caaacatcca tttctgtacc aagaccctca 954  
 ctcgaatgtc actatcccaa gatgaaacct aaaaatcact ttccattctt tcttgatctt 1014  
 accccaccat ccaatcagct gccatgccca gtttagttaa ccccccaa gctgcttcat 1074  
 gcaaccttcc attcctattc cttttgccaa cccatgatgt agagatgtgg attcatgaca 1134  
 ttttgttcat acaacttctt caataaaaca ttataatatg tgccccaaag ataaagctga 1194  
 agaatgagat gaatgtgaaa ttaaagggtt gcattgtcttc ctaatcctaa 1244

<210> 2

<211> 157

<212> PRT

<213> Homo sapiens

<400> 2

Met Asn Pro Gln Arg Glu Ala Ala Pro Lys Ser Tyr Ala Ile Arg Asp  
 1 5 10 15

Ser Arg Gln Met Val Trp Val Leu Ser Gly Asn Ser Leu Ile Ala Ala  
 20 25 30

Pro Leu Ser Arg Ser Ile Lys Pro Val Thr Leu His Leu Ile Ala Cys  
 35 40 45

Arg Asp Thr Glu Phe Ser Asp Lys Glu Lys Gly Asn Met Val Tyr Leu

50                      55                      60  
 Gly Ile Lys Gly Lys Asp Leu Cys Leu Phe Cys Ala Glu Ile Gln Gly  
 65                      70                      75                      80  
 Lys Pro Thr Leu Gln Leu Lys Glu Lys Asn Ile Met Asp Leu Tyr Val  
                     85                      90                      95  
 Glu Lys Lys Ala Gln Lys Pro Phe Leu Phe Phe His Asn Lys Glu Gly  
                     100                      105                      110  
 Ser Thr Ser Val Phe Gln Ser Val Ser Tyr Pro Gly Trp Phe Ile Ala  
                     115                      120                      125  
 Thr Ser Thr Thr Ser Gly Gln Pro Ile Phe Leu Thr Lys Glu Arg Gly  
                     130                      135                      140  
 Ile Thr Asn Asn Thr Asn Phe Tyr Leu Asp Ser Val Glu  
 145                      150                      155

<210> 3  
 <211> 164  
 <212> PRT  
 <213> Homo sapiens

<400> 3  
 Met Asn Pro Gln Arg Glu Ala Ala Pro Lys Ser Tyr Ala Ile Arg Asp  
 1                      5                      10                      15  
 Ser Arg Gln Met Val Trp Val Leu Ser Gly Asn Ser Leu Ile Ala Ala  
                     20                      25                      30  
 Pro Leu Ser Arg Ser Ile Lys Pro Val Thr Leu His Leu Ile Ala Cys  
                     35                      40                      45  
 Arg Asp Thr Glu Phe Ser Asp Lys Glu Lys Gly Asn Met Val Tyr Leu  
                     50                      55                      60  
 Gly Ile Lys Gly Lys Asp Leu Cys Leu Phe Cys Ala Glu Ile Gln Gly  
 65                      70                      75                      80  
 Lys Pro Thr Leu Gln Leu Lys Leu Gln Gly Ser Gln Asp Asn Ile Gly  
                     85                      90                      95  
 Lys Asp Thr Cys Trp Lys Leu Val Gly Ile His Thr Cys Ile Asn Leu  
                     100                      105                      110  
 Asp Val Arg Glu Ser Cys Phe Met Gly Thr Leu Asp Gln Trp Gly Ile  
                     115                      120                      125  
 Gly Val Gly Arg Lys Lys Trp Lys Ser Ser Phe Gln His His His Leu  
                     130                      135                      140  
 Arg Lys Lys Asp Lys Asp Phe Ser Ser Met Arg Thr Asn Ile Gly Met  
 145                      150                      155                      160

Pro Gly Arg Met

<210> 4  
<211> 158  
<212> PRT  
<213> Homo sapiens

<400> 4  
Met Glu Lys Ala Leu Lys Ile Asp Thr Pro Gln Gln Gly Ser Ile Gln  
1 5 10 15  
Asp Ile Asn His Arg Val Trp Val Leu Gln Asp Gln Thr Leu Ile Ala  
20 25 30  
Val Pro Arg Lys Asp Arg Met Ser Pro Val Thr Ile Ala Leu Ile Ser  
35 40 45  
Cys Arg His Val Glu Thr Leu Glu Lys Asp Arg Gly Asn Pro Ile Tyr  
50 55 60  
Leu Gly Leu Asn Gly Leu Asn Leu Cys Leu Met Cys Ala Lys Val Gly  
65 70 75 80  
Asp Gln Pro Thr Leu Gln Leu Lys Glu Lys Asp Ile Met Asp Leu Tyr  
85 90 95  
Asn Gln Pro Glu Pro Val Lys Ser Phe Leu Phe Tyr His Ser Gln Ser  
100 105 110  
Gly Arg Asn Ser Thr Phe Glu Ser Val Ala Phe Pro Gly Trp Phe Ile  
115 120 125  
Ala Val Ser Ser Glu Gly Gly Cys Pro Leu Ile Leu Thr Gln Glu Leu  
130 135 140  
Gly Lys Ala Asn Thr Thr Asp Phe Gly Leu Thr Met Leu Phe  
145 150 155

<210> 5  
<211> 177  
<212> PRT  
<213> Homo sapiens

<400> 5  
Met Glu Ile Cys Arg Gly Leu Arg Ser His Leu Ile Thr Leu Leu Leu  
1 5 10 15  
Phe Leu Phe His Ser Glu Thr Ile Cys Arg Pro Ser Gly Arg Lys Ser  
20 25 30  
Ser Lys Met Gln Ala Phe Arg Ile Trp Asp Val Asn Gln Lys Thr Phe  
35 40 45  
Tyr Leu Arg Asn Asn Gln Leu Val Ala Gly Tyr Leu Gln Gly Pro Asn

50	55	60
Val Asn Leu Glu Glu Lys Ile Asp Val Val Pro Ile Glu Pro His Ala		
65	70	75 80
Leu Phe Leu Gly Ile His Gly Gly Lys Met Cys Leu Ser Cys Val Lys		
	85	90 95
Ser Gly Asp Glu Thr Arg Leu Gln Leu Glu Ala Val Asn Ile Thr Asp		
	100	105 110
Leu Ser Glu Asn Arg Lys Gln Asp Lys Arg Phe Ala Phe Ile Arg Ser		
	115	120 125
Asp Ser Gly Pro Thr Thr Ser Phe Glu Ser Ala Ala Cys Pro Gly Trp		
	130	135 140
Phe Leu Cys Thr Ala Met Glu Ala Asp Gln Pro Val Ser Leu Thr Asn		
145	150	155 160
Met Pro Asp Glu Gly Val Met Val Thr Lys Phe Tyr Phe Gln Glu Asp		
	165	170 175

Glu

<210> 6  
 <211> 153  
 <212> PRT  
 <213> Homo sapiens

<400> 6

Ala Pro Val Arg Ser Leu Asn Cys Thr Leu Arg Asp Ser Gln Gln Lys		
1	5	10 15
Ser Leu Val Met Ser Gly Pro Tyr Glu Leu Lys Ala Leu His Leu Gln		
	20	25 30
Gly Gln Asp Met Glu Gln Gln Val Val Phe Ser Met Ser Phe Val Gln		
	35	40 45
Gly Glu Glu Ser Asn Asp Lys Ile Pro Val Ala Leu Gly Leu Lys Glu		
50	55	60
Lys Asn Leu Tyr Leu Ser Cys Val Leu Lys Asp Asp Lys Pro Thr Leu		
65	70	75 80
Gln Leu Glu Ser Val Asp Pro Lys Asn Tyr Pro Lys Lys Lys Met Glu		
	85	90 95
Lys Arg Phe Val Phe Asn Lys Ile Glu Ile Asn Asn Lys Leu Glu Phe		
	100	105 110
Glu Ser Ala Gln Phe Pro Asn Trp Tyr Ile Ser Thr Ser Gln Ala Glu		
	115	120 125

Asn Met Pro Val Phe Leu Gly Gly Thr Lys Gly Gly Gln Asp Ile Thr  
 130 135 140

Asp Phe Thr Met Gln Phe Val Ser Ser  
 145 150

<210> 7  
 <211> 11  
 <212> PRT  
 <213> Human immunodeficiency virus type 1

<400> 7  
 Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg  
 1 5 10

<210> 8  
 <211> 15  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: internalizing  
 domain derived from HIV tat protein

<400> 8  
 Gly Gly Gly Gly Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg  
 1 5 10 15

<210> 9  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide 2362-94

<400> 9  
 catggacctg tatgtggaga aga 23

<210> 10  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide 2362-95

<400> 10  
 gccagggtaa gagactgact gaa 23

<210> 11  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide 870-02

<400> 11  
agcggataac aatttcacac agg

23

<210> 12  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide 2366-21

<400> 12  
gcctaggctg gatttattcc acag

24

<210> 13  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide 1019-06

<400> 13  
gctctaatac gactcactat aggg

24

<210> 14  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide 2362-98

<400> 14  
ctgatgtggt ggaggtggct at

22

<210> 15  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
     Oligonucleotide 1340-35  
  
 <400> 15  
 cccagtcacg acgttgtaaa acg 23  
  
  
 <210> 16  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence:  
     Oligonucleotide 1019-05  
  
 <400> 16  
 tgaatttagg tgacactata gaagag 26  
  
  
 <210> 17  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence:  
     Oligonucleotide 2379-15  
  
 <400> 17  
 gtccctccaga gcctcaagag atc 23  
  
  
 <210> 18  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence:  
     Oligonucleotide 2375-10  
  
 <400> 18  
 ttaggattag gaagacatgc aaacc 25